

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 September 2002 (06.09.2002)

PCT

(10) International Publication Number
WO 02/068595 A2

(51) International Patent Classification⁷: C12N

(21) International Application Number: PCT/US02/05474

(22) International Filing Date: 21 February 2002 (21.02.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/271,307	23 February 2001 (23.02.2001)	US
09/794,715	26 February 2001 (26.02.2001)	US
10/046,924	14 January 2002 (14.01.2002)	US

(71) Applicant (for all designated States except US): **THE REGENTS OF THE UNIVERSITY OF CALIFORNIA** [US/US]; 1111 Franklin Street, 5th Floor, Oakland, CA 94607 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CASES, Sylvaine** [FR/US]; 9 Meadow Park, Belmont, CA 94002 (US).

STONE, Scot [CA/US]; 2550 Hilborn Dr., #168, Fairfield, CA 94533 (US). **ZHOU, Ping** [CN/US]; 886 Stonehaven Drive, Walnut Creek, CA 94598 (US). **FARESE, Robert, V.** [US/US]; 25 6th Avenue, San Francisco, CA 94118 (US). **YEN, Chi-Liang, Eric** [CN/US]; 1427 22nd Avenue, San Francisco, CA 94122 (US).


(74) Agent: **BORDEN, Paula, A.**; BOZICEVIC, FIELD & FRANCIS, L.L.P., 200 Middlefield Road, Suite 200, Menlo Park, CA 94025 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,

[Continued on next page]

(54) Title: MONO-AND DIACYGLYCEROL ACYLTRANSFERASES AND METHODS OF USE THEREOF



(57) Abstract: Nucleic acid compositions encoding polypeptide products with diglyceride acyltransferase and/or monoacylglycerol acyltransferase activity, as well as the polypeptide products encoded thereby, i.e., mammalian DGAT2 and MGAT1 polypeptide products, and methods for producing the same, are provided. Also provided are: methods and compositions for modulating DGAT2 and MGAT1 activity; DGAT2 and MGAT1 transgenic cells, animals and plants, as well as methods for their preparation; and methods for making diglyceride, diglyceride compositions, triglycerides and triglyceride compositions, as well as the compositions produced by these methods. The subject methods and compositions find use in a variety of different applications, including research, medicine, agriculture and industry applications.

WO 02/068595 A2



GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

- *without international search report and to be republished upon receipt of that report*

MONO- AND DIACYLGLYCEROL ACYLTRANSFERASES AND METHODS OF USE THEREOF**FIELD OF THE INVENTION**

The field of the invention is enzymes, particularly acyltransferases.

BACKGROUND OF THE INVENTION

5 Diacylglycerol O-Acyltransferase (EC 2.3.1.20), also known as diglyceride acyltransferase or DGAT, is a critical enzyme in triacylglycerol synthesis. Triacylglycerols are quantitatively the most important storage form of energy for eukaryotic cells. DGAT catalyzes the rate-limiting and terminal step in triacylglycerol synthesis using diacylglycerol and fatty acyl CoA as substrates. As such, DGAT plays a fundamental role in the metabolism of cellular diacylglycerol and is important in higher eukaryotes for intestinal fat absorption, lipoprotein assembly, fat storage in adipocytes, milk production and possibly egg production and sperm maturation.

15 Diacylglycerol is the precursor of such important lipids as triacylglycerol and phospholipids, which store energy and form cellular membranes. In eukaryotes, two major pathways for synthesizing diacylglycerol exist: the glycerol phosphate pathway and the monoacylglycerol pathway. Both pathways generate diacylglycerol that can be used as a substrate by acyl CoA:diacylglycerol acyltransferase (DGAT) for triacylglycerol synthesis. In the glycerol phosphate pathway, which functions in most cells, diacylglycerol is derived by the dephosphorylation of phosphatidic acid produced by sequential acylations of glycerol phosphate. In the monoacylglycerol pathway, which has been reported predominantly in the intestine, diacylglycerol is formed directly from monoacylglycerol and fatty acyl CoA in a reaction catalyzed by monoacylglycerol acyltransferase (MGAT) (E.C. 2.3.1.22).

25 MGAT is best known for its role in fat absorption in the intestine, where the fatty acids and *sn*-2-monoacylglycerol generated from the digestion of dietary fat (mainly triacylglycerol) are resynthesized into triacylglycerol in enterocytes for chylomicron synthesis and secretion. MGAT catalyzes the first step of this process, in which fatty acyl CoA, formed from fatty acids and CoA, and *sn*-2-monoacylglycerol are covalently joined. Because the monoacylglycerol pathway predominates in intestinal triacylglycerol synthesis, MGAT may be a pharmaceutical target for modulating fat absorption.

30 MGAT activity is also found at high levels in liver of suckling rats and in white adipose tissue of migrating sparrows, where triacylglycerols are actively hydrolyzed to

provide fatty acids for energy. MGAT preferentially acylates monoacylglycerols that contain a polyunsaturated fatty acyl moiety at the *sn*-2 position. Thus, MGAT may preserve essential fatty acids, all of which are polyunsaturated, by resynthesizing them into triacylglycerols.

This function may be relevant in mammalian white adipose tissue, which possesses

5 significant levels of MGAT activity. In addition, MGAT may also play a role in signaling, since its product, diacylglycerol, and one of its substrates, 2-arachidonoylglycerol, are signaling molecules.

Like many enzymes that participate in neutral lipid synthesis, MGAT has proven difficult to purify to homogeneity, and an MGAT gene has not been identified. Several
10 partial purifications of MGAT enzymes have been reported, and a 43-kDa MGAT enzyme was purified recently from peanut cotyledons. Difficulties in the purification of MGAT may reflect its hydrophobicity or its involvement in an enzyme complex.

Because of its central role in a variety of different processes, there is much interest in the identification of polynucleotides encoding proteins having DGAT and MGAT activity,
15 as well as the proteins encoded thereby.

Relevant Literature

Of particular interest are: U.S. Patent No. 6,100,077; and PCT Published Application Nos. WO 98/55631; WO 99/67268; WO 00/01713; WO 99/67403; WO 00/32793; WO
20 00/32756; WO 00/36114; WO 00/60095; WO 00/66749.

Also of interest are: Smith et al., *Nat. Genet.* 2000 (25), 87-90). Cases et al., *Proc. Natl. Acad. Sci. USA* (October 1998) 95:13018-13023; Oelkers et al., *J. Biol. Chem.* (October 9, 1998) 273:26765-71; and Cases et al. (2001) *J. Biol. Chem.* 276:38870-38876.

References describing the role DGAT plays in various biological processes include:
25 Bell & Coleman, *Annu. Rev. Biochem.* (1980) 49: 459-487; Lehner & Kuksis, *Prog. Lipid Res.* (1996) 35: 169-201; Brindley, *Biochemistry of Lipids, Lipoproteins and Membranes* (eds. Vance & Vance)(Elsevier, Amsterdam)(1991) pp171-203; Haagsman & Van Golde, *Arch. Biochem. Biophys.* (1981) 208:395-402; Coleman & Bell, *J. Biol. Chem.* (1976) 251:4537-4543.

30 References discussing MGAT activity and purification include: Coleman and Haynes (1984) *J. Biol. Chem.* 259:8934-8938; Mostafa et al. (1994) *Lipids* 29:785-791; Xia et al. 1993) *Am. J. Physiol.* 265:R414-R419; Jamdar et al. (1992) *Arch. Biochem. Biophys.* 296:419-425; Manganaro et al. (1985) *Can. J. Biochem. Cell Biol.* 63:341-347; Bhat et al.

(1993) *Arch. Biochem. Biophys.* 300:663-669; Tumaney et al. (2001) *J. Biol. Chem.* 276:10847-10852; Lehner and Kuksis (1995) *J. Biol. Chem.* 270:13630-13636.

SUMMARY OF THE INVENTION

5 Nucleic acid compositions encoding polypeptide products with diglyceride acyltransferase and/or monoacylglycerol acyltransferase activity, as well as the polypeptide products encoded thereby, i.e., mammalian DGAT2 α and MGAT1 polypeptide products, and methods for producing the same, are provided. Also provided are: methods and compositions for modulating DGAT2 α and MGAT1 activity; DGAT2 α and MGAT1
10 transgenic cells, animals and plants, as well as methods for their preparation; and methods for making diglyceride, diglyceride compositions, triglycerides and triglyceride compositions, as well as the compositions produced by these methods. The subject methods and compositions find use in a variety of different applications, including research, medicine, agriculture and industry applications.

BRIEF DESCRIPTION OF THE FIGURES

Fig. 1 provides a graphical representation of the results obtained from a pulse assay that demonstrates the existence of mouse DGAT2 α .

Fig. 2 provides a hydrophobicity plot of mouse DGAT2 α .

20 Figs 3A to 3C provide graphical results of various mouse DGAT2 α activity assays.

Fig. 4 provides the expression profile for mouse DGAT2 α .

Fig. 5 provides the results of an assay showing that mouse DGAT2 α expression increases during 3T3-L1 adipocyte differentiation.

25 Figs 6A and 6B provide the amino acid and nucleic acid sequences of mouse DGAT2 α .

Figs. 7A and 7B provide the amino acid and nucleic acid sequences of human DGAT2 α .

Figures 8A-D provide the amino acid and nucleic acid sequences of various mouse and human DGAT2 α homologs.

30 Figure 9 depicts schematically two major pathways for synthesizing diacylglycerol (DAG).

Figure 10A depicts a comparison of the amino acid sequences of mouse MGAT1 and mouse DGAT2. Figure 10B depicts a hydrophobicity plot of mouse MGAT1.

Figure 11 depicts graphs showing MGAT1 activity using oleoyl CoA or 2-monooleoylglycerol.

Figure 12 depicts graphs showing that MGAT1 has activity toward all stereoisomers of monoacyl glycerol.

5 Figure 13 depicts expression of MGAT1 protein in COS-7 cells.

Figure 14 depicts the tissue distribution of MGAT1 mRNA expression.

Figures 15 A-D depict amino acid sequences of human and mouse DGAT/MGAT enzymes.

10

DETAILED DESCRIPTION OF THE INVENTION

Nucleic acid compositions encoding polypeptide products with diglyceride and/or monoacylglycerol acyltransferase activity, as well as the polypeptide products encoded thereby, i.e., mammalian DGAT2 α and MGAT1, and methods for producing the same, are provided. In many embodiments, the subject nucleic acids encode enzymes that exhibit
15 monoacylglycerol acyltransferase activity, diacylglycerol acyltransferase activity, or both mono- and diacyltransferase activity. For example, DGAT2 α polypeptides exhibit diglyceride acyltransferase activity (also referred to herein as "DGAT2" polypeptides); and mammalian MGAT1 polypeptides (also referred to herein as "DC2" polypeptides) exhibit monoacylglycerol acyltransferase activity, and in some embodiments also exhibit
20 diacylglycerol acyltransferase activity.

Also provided are: methods and compositions for modulating DGAT2 α and MGAT1 activity, e.g. in the treatment of disease conditions associated with DGAT2 α and/or MGAT1 activity, including obesity; MGAT1 and DGAT2 α transgenic cells, animals, plants and fungi, and methods for their preparation, e.g. for use in research, food production, industrial
25 feedstock production, etc.; and methods for making diglycerides, diglyceride compositions, triglycerides, and triglyceride compositions, e.g. oils. The methods and compositions of the subject invention find use in a variety of different applications and fields, including research, medicine, agriculture and industry.

30

Before the subject invention is further described, it is to be understood that the invention is not limited to the particular embodiments of the invention described below, as variations of the particular embodiments may be made and still fall within the scope of the appended claims. It is also to be understood that the terminology employed is for the purpose

of describing particular embodiments, and is not intended to be limiting. Instead, the scope of the present invention will be established by the appended claims.

In this specification and the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention belongs.

NUCLEIC ACID COMPOSITIONS

Nucleic acid compositions encoding polypeptide products, as well as fragments thereof, having mono- and/or diglyceride acetyltransferase activity are provided. In many embodiments, the subject nucleic acids encode enzymes that exhibit monoacylglycerol acyltransferase activity, diacylglycerol acyltransferase activity, or both mono- and diacyltransferase activity. Specifically, nucleic acid compositions encoding mammalian, e.g., human, mouse, etc., DGAT2 α polypeptides having diglyceride acyltransferase activity (also referred to herein as "DGAT2" polypeptides), and mammalian MGAT1 polypeptides exhibiting monoacylglycerol acyltransferase activity (also referred to herein as "DC2" polypeptides), are provided.

By nucleic acid composition is meant a composition comprising a sequence of DNA having an open reading frame that encodes a DGAT2 α or an MGAT1 polypeptide, i.e. a gene or genomic region encoding a polypeptide having mono- and/or diglyceride acyltransferase activity, and is capable, under appropriate conditions, of being expressed as a DGAT2 α or an MGAT1 polypeptide.

Also encompassed in this term are nucleic acids that are homologous or substantially similar or identical to the nucleic acids encoding DGAT2 α polypeptides, or MGAT1 polypeptides. Thus, the subject invention provides nucleic acids encoding mammalian DGAT2 α , such as nucleic acids encoding human DGAT2 α and homologs thereof and mouse DGAT2 α and homologs thereof. The subject invention provides nucleic acids encoding mammalian MGAT1, such as nucleic acids encoding mouse MGAT1 (also referred to herein as "DC2"), and homologs thereof.

The coding sequence of the human DGAT2 α genomic sequence, i.e. the human cDNA encoding the human DGAT2 α enzyme, includes or comprises a nucleic acid

sequence substantially the same as or identical to that identified as SEQ ID NO:01 or SEQ ID NO:18, *infra*. The coding sequence of the mouse DGAT2 α genomic sequence, i.e., the mouse cDNA encoding the mouse DGAT2 α enzyme, includes or comprises a nucleic acid substantially the same as or identical to the sequence identified as SEQ ID NO:03, *infra*.

- 5 The coding sequence of the mouse MGAT1 genomic sequence, i.e. the mouse cDNA encoding the mouse MGAT1 enzyme, includes or comprises a nucleic acid sequence substantially the same as or identical to that identified as SEQ ID NO:05, *infra*.

The source of homologous nucleic acids to those specifically listed above may be any species, including both animal and plant species, e.g., primate species, particularly human; rodents, such as rats and mice, canines, felines, bovines, ovines, equines, yeast, nematodes, etc. Between mammalian species, e.g., human and mouse, homologs have substantial sequence similarity, e.g. at least 75% sequence identity, usually at least 90%, more usually at least 95% between nucleotide sequences. Sequence similarity is calculated based on a reference sequence, which may be a subset of a larger sequence, such as a conserved motif, coding region, flanking region, etc. A reference sequence will usually be at least about 18 nt long, more usually at least about 30 nt long, and may extend to the complete sequence that is being compared. Algorithms for sequence analysis are known in the art, such as BLAST, described in Altschul *et al.* (1990), *J. Mol. Biol.* **215**:403-10. Unless specified otherwise, all sequence identity values provided herein are determined using GCG (Genetics Computer Group, Wisconsin Package, Standard Settings, gap creation penalty 3.0, gap extension penalty 0.1). The sequences provided herein are essential for recognizing DGAT2 α - related and homologous polynucleotides in database searches. Specific DGAT2 α homologues of interest are provide in Fig. 8, i.e., SEQ ID NOs. 05, 07, 09, 11, 13 and 15; and in Figure 15, i.e., SEQ ID NOs: 19, 20, 21, and 22.

- 25 Also provided are nucleic acids that hybridize to the above-described specific nucleic acids, e.g., those nucleic acids having a sequence of SEQ ID NO:01, 03, 05, 07, 09, 11, 13, 15, or 18, or a coding sequence of any one of the foregoing sequences, under stringent conditions. An example of stringent hybridization conditions is hybridization at 50°C or higher and 0.1 \times SSC (15 mM sodium chloride/1.5 mM sodium citrate). Another example of stringent hybridization conditions is overnight incubation at 42°C in a solution: 50% formamide, 5 \times SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5 \times Denhardt's solution, 10% dextran sulfate, and 20 μ g/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1 \times SSC at about 65°C. Stringent
- 30

hybridization conditions are hybridization conditions that are at least as stringent as the above representative conditions. Other stringent hybridization conditions are known in the art and may also be employed to identify nucleic acids of this particular embodiment of the invention.

Also provided are nucleic acids that encode a polypeptide having mono and/or diacylglycerol acyltransferase activity and having at least about 50%, at least about 60%, at least about 70%, at least about 75%, at least about 80%, at least about 90%, at least about 95%, or higher, nucleotide sequence identity to a nucleic acid having a nucleic acid sequence set forth in any one of SEQ ID NO:01, 03, 05, 07, 09, 11, 13, 15, or 18. Also provided are nucleic acids that encode a polypeptide having mono- and/or diacylglycerol acyltransferase activity and having at least about 50%, at least about 60%, at least about 70%, at least about 75%, at least about 80%, at least about 90%, at least about 95%, or higher, nucleotide sequence identity to the coding region of a nucleic acid having a sequence set forth in any one of SEQ ID NO:01, 03, 05, 07, 09, 11, 13, 15, or 18.

Nucleic acids encoding the DGAT2 α proteins, DGAT2 α polypeptides, and MGAT polypeptides of the subject invention may be cDNAs or genomic DNAs, i.e. portions of chromosomes that include both introns and exons, as well as promoter regions, etc., as well as fragments thereof. The term "DGAT2 α - gene" shall be intended to mean the open reading frame encoding specific DGAT2 α proteins and polypeptides, and DGAT2 α introns, as well as adjacent 5' and 3' non-coding nucleotide sequences involved in the regulation of expression, up to about 20 kb beyond the coding region, but possibly further in either direction. Similarly, the term "MGAT1 gene" refers to the open reading from encoding specific MGAT polypeptides, and MGAT1 introns, as well as adjacent 5' and 3' non-coding nucleotide sequences involved in the regulation of expression, up to about 20 kb beyond the coding region, but possibly further in either direction. The gene may be introduced into an appropriate vector for extrachromosomal maintenance or for integration into a host genome.

The term "cDNA" as used herein is intended to include all nucleic acids that share the arrangement of sequence elements found in native mature mRNA species, where sequence elements are exons and 3' and 5' non-coding regions. Normally mRNA species have contiguous exons, with the intervening introns, when present, being removed by nuclear RNA splicing, to create a continuous open reading frame encoding an MGAT1 or a DGAT2 α protein.

Also provided are nucleic acids that encode the DGAT2 α and MGAT1 proteins encoded by the above-described nucleic acids, but differ in sequence from the above-described nucleic acids due to the degeneracy of the genetic code. Also provided are nucleic acids that encode DGAT2 α and MGAT1 proteins that include conservative amino acid changes when compared to, e.g., the amino acid sequences set forth in SEQ ID NO:02 or 06.

A genomic sequence of interest comprises the nucleic acid present between the initiation codon and the stop codon, as defined in the listed sequences, including all of the introns that are normally present in a native chromosome. It may further include the 3' and 5' untranslated regions found in the mature mRNA. It may further include specific transcriptional and translational regulatory sequences, such as promoters, enhancers, *etc.*, including about 1 kb, but possibly more, of flanking genomic DNA at either the 5' or 3' end of the transcribed region. The genomic DNA may be isolated as a fragment of 100 kbp or smaller; and substantially free of flanking chromosomal sequence. The genomic DNA flanking the coding region, either 3' or 5', or internal regulatory sequences as sometimes found in introns, contains sequences required for proper tissue and stage specific expression.

The nucleic acid compositions of the subject invention may encode all or a part of the subject DGAT2 α or MGAT1 proteins and polypeptides, described in greater detail *infra*. Double or single stranded fragments may be obtained from the DNA sequence by chemically synthesizing oligonucleotides in accordance with conventional methods, by restriction enzyme digestion, by PCR amplification, *etc.* For the most part, DNA fragments will be of at least 15 nt, usually at least 18 nt or 25 nt, and may be at least about 50 nt.

The MGAT1 and DGAT2 α - nucleic acids or genes of the subject invention are isolated and obtained in substantial purity, generally as other than an intact chromosome. Usually, the DNA will be obtained substantially free of other nucleic acid sequences that do not include a DGAT2 α sequence or fragment thereof, generally being at least about 50%, usually at least about 90% pure and are typically "recombinant", *i.e.* flanked by one or more nucleotides with which it is not normally associated on a naturally occurring chromosome.

In addition to the plurality of uses described in greater detail in following sections, the subject nucleic acid compositions find use in the preparation of all or a portion of the DGAT2 α or MGAT1 polypeptides, as described below.

POLYPEPTIDE COMPOSITIONS

Also provided by the subject invention are polypeptides having mono- and/or diglyceride acyltransferase activity, *i.e.*, capable of catalyzing the acylation of

diacylglycerol, acylation of monoacylglycerol, or acylation of both mono- and diacylglycerol. Such enzymes are referred to herein as "mono- and diacylglycerol acyltransferases." Examples of such polypeptides are DGAT2 α (also referred to as "DGAT2") and MGAT1 (also referred to as "DC2"). The term "polypeptide composition" as used herein refers to both full-length proteins as well as portions or fragments thereof. Also included in this term are variations of the naturally occurring proteins, where such variations are homologous or substantially similar to the naturally occurring protein, as described in greater detail below, be the naturally occurring protein the human protein, mouse protein, or protein from some other mammalian species which naturally expresses a subject acyltransferase. In the following description of the subject invention, the term "DGAT2 α " is used to refer not only to the human form of the enzyme, but also to homologs thereof expressed in non-human mammalian species. Similarly, the term "mouse MGAT1" refers not only to the mouse form of the enzyme, but also to homologs thereof expressed in other mammalian species.

The subject mono- and diacylglycerol acyltransferases are, in their natural environment, trans-membrane proteins. The subject proteins are characterized by the presence of at least one potential N-linked glycosylation site, at least one potential tyrosine phosphorylation site, and multiple hydrophobic domains, including 4 to 12, e.g., 6, hydrophobic domains capable of serving as trans-membrane regions. The proteins range in length from about 300 to 500, usually from about 325 to 475 and more usually from about 350 to 425 amino acid residues, and the projected molecular weight of the subject proteins based solely on the number of amino acid residues in the protein ranges from about 35 to 55, usually from about 37.5 to 47.5 and more usually from about 40 to 45 kDa, where the actual molecular weight may vary depending on the amount of glycosylation of the protein and the apparent molecular weight may be considerably less because of SDS binding on gels.

The amino acid sequences of the subject proteins are characterized by having substantially no homology to the known DGAT enzymes. More specifically, the subject human DGAT2 α and mouse MGAT1 enzymes have substantially no homology to the human DGAT enzyme described in Cases et al., "Identification of a gene encoding an acyl CoA:diacylglycerol acyltransferase, a key enzyme in triacylglycerol synthesis," Proc. Natl. Acad. Sci. U.S.A. 95 (22), 13018-13023 (1998). Likewise, the subject mouse DGAT2 α enzymes have substantially no homology to the mouse DGAT enzyme described in Cases et al., "Identification of a gene encoding an acyl CoA:diacylglycerol acyltransferase, a key

enzyme in triacylglycerol synthesis," Proc. Natl. Acad. Sci. U.S.A. 95 (22), 13018-13023 (1998). By substantially no homology is meant that the homology does not exceed about 20%, and usually will not exceed about 10% and more usually will not exceed about as determined using GCG (Genetics Computer Group, Wisconsin Package, Standard Settings, Gap Creation Penalty 3.0, Gap Extension Penalty 0.1).

Of particular interest in many embodiments are proteins that are non-naturally glycosylated. By non-naturally glycosylated is meant that the protein has a glycosylation pattern, if present, which is not the same as the glycosylation pattern found in the corresponding naturally occurring protein. For example, human DGAT2 α of the subject invention and of this particular embodiment is characterized by having a glycosylation pattern, if it is glycosylated at all, that differs from that of naturally occurring human DGAT2 α . Thus, the non-naturally glycosylated DGAT2 α proteins of this embodiment include non-glycosylated DGAT2 α proteins, i.e. proteins having no covalently bound glycosyl groups.

The sequence of the full-length human DGAT2 α protein is identified, *infra*, as SEQ ID NO:02. As such, DGAT2 α proteins having an amino acid sequence that is substantially the same as or identical to the sequence of SEQ ID NO:2 are of interest. By substantially the same as is meant a protein having a region with a sequence that has at least about 75%, usually at least about 90% and more usually at least about 98 % sequence identity with the sequence of SEQ ID NO:02, as measured by GCG, *supra*. Of particular interest in other embodiments is the mouse DGAT2 α protein, where the mouse DGAT2 α protein of the subject invention has an amino acid sequence that is substantially the same as or identical to the sequence appearing as SEQ ID NO:04, *infra*.

In addition to the specific mammalian DGAT2 α proteins described above, homologs or proteins (or fragments thereof) from other species, i.e. other animal or plant species, are also provided, where such homologs or proteins may be from a variety of different types of species, including animals, such as mammals, e.g., rodents, such as rats, mice; domestic animals, e.g. horse, cow, dog, cat; humans, and the like. By homolog is meant a protein having at least about 35 %, usually at least about 40% and more usually at least about 60 % amino acid sequence identity the specific DGAT2 α proteins as identified in SEQ ID NOS: 02 to 04, where sequence identity is determined using GCG, *supra*. Specific homologs of interest include human DC 2, human DC3, human DC4, human DC5, mouse DC2 and mouse DC3, the sequences of which are provided in Fig. 8 (i.e., SEQ ID NOs. 06, 08, 10, 12,

14 and 16); and human and mouse variants of DC5, which are shown in Figure 15 (i.e., SEQ ID NOs: 19, 20, 21, and 22).

Mouse MGAT1 exhibits monoacylglycerol acyltransferase activity. The sequence of the full-length mouse MGAT1 protein is identified, *infra*, as SEQ ID NO:06 (identified as
5 “mouse DC2” in Figure 8). As such, subject MGAT1 proteins having an amino acid sequence that is substantially the same as or identical to the sequence of SEQ ID NO:06 are of interest. By substantially the same as is meant a protein having a region with a sequence that has at least about 75%, usually at least about 90% and more usually at least about 98 % sequence identity with the sequence of SED ID NO:06, as measured by GCG, *supra*.

10 Mono- and diacylglycerol acyltransferases of the subject invention (e.g. human DGAT2 α or a homolog thereof; non-human DGAT2 α proteins, e.g. mouse DGAT2 α ; mouse MGAT1 polypeptide or a homolog thereof) are present in a non-naturally occurring environment, *e.g.* are separated from their naturally occurring environment. In certain
15 embodiments, the subject mono- and diacylglycerol acyltransferases are present in a composition that is enriched for such an enzyme, *e.g.*, enriched for DGAT2 α as compared to DGAT2 α in its naturally occurring environment. As such, purified mono- and diacylglycerol acyltransferases are provided, where by purified is meant that subject enzyme is present in a composition that is substantially free of proteins other than the subject enzyme, where by
20 substantially free is meant that less than 90 %, usually less than 60 % and more usually less than 50 % of the composition is made up of proteins other than the subject enzyme. For example, for compositions that are enriched for DGAT2 α proteins, such compositions will exhibit a DGAT2 α activity of at least about 100, usually at least about 200 and more usually at least about 1000 pmol triglycerides formed/mg protein/min, where such activity is determined by the assay described in the Experimental Section, *infra*.

25 In certain embodiments of interest, a subject enzyme is present in a composition that is substantially free of the constituents that are present in its naturally occurring environment. For example, a human DGAT2 α protein comprising composition according to the subject invention in this embodiment will be substantially, if not completely, free of those other biological constituents, such as proteins, carbohydrates, lipids, etc., with which it
30 is present in its natural environment. As such, protein compositions of these embodiments will necessarily differ from those that are prepared by purifying the protein from a naturally occurring source, where at least trace amounts of the protein's constituents will still be present in the composition prepared from the naturally occurring source.

The mono- and diacylglycerol acyltransferases of the subject invention may also be present as an isolate, by which is meant that the subject enzyme is substantially free of both proteins other than a subject enzyme and other naturally occurring biologic molecules, such as oligosaccharides, polynucleotides and fragments thereof, and the like, where substantially free in this instance means that less than 70 %, usually less than 60% and more usually less than 50 % (dry weight) of the composition containing the isolated subject enzyme is a naturally occurring biological molecule other than the subject enzyme. In certain embodiments, the subject enzyme is present in substantially pure form, where by substantially pure form is meant at least 95%, usually at least 97% and more usually at least 99% pure.

In addition to the naturally occurring subject proteins, mono- and diacylglycerol acyltransferase polypeptides which vary from the naturally occurring DGAT2 α and/or MGAT1 proteins are also provided. By "DGAT2 α polypeptides" and "MGAT1 polypeptides" is meant proteins having an amino acid sequence encoded by an open reading frame (ORF) of a DGAT2 α gene or an MGAT1 gene, respectively, as described *supra*, including the full length DGAT2 α or MGAT1 protein and fragments thereof, particularly biologically active fragments and/or fragments corresponding to functional domains; and including fusions of the subject polypeptides to other proteins or parts thereof. Fragments of interest will typically be at least about 10 amino acids (aa) in length, usually at least about 50 aa in length, and may be as long as 300 aa in length or longer, but will usually not exceed about 1000 aa in length, where the fragment will have a stretch of amino acids that is identical to a subject protein of SEQ ID NO:2, SEQ ID NO:04, or SEQ ID NO:06 or a homolog thereof; of at least about 10 aa, and usually at least about 15 aa, and in many embodiments at least about 50 aa in length.

PREPARATION OF SUBJECT POLYPEPTIDES

The subject proteins and polypeptides may be obtained from naturally occurring sources, but are preferably synthetically produced. Where obtained from naturally occurring sources, the source chosen will generally depend on the species from which the subject protein is to be derived.

The subject polypeptide compositions may be synthetically derived by expressing a recombinant gene encoding the subject protein, such as the polynucleotide compositions described above, in a suitable host. For expression, an expression cassette may be employed.

The expression vector will provide a transcriptional and translational initiation region, which may be inducible or constitutive, where the coding region is operably linked under the transcriptional control of the transcriptional initiation region, and a transcriptional and translational termination region. These control regions may be native to a DGAT2 α gene, an MGAT1 gene, or may be derived from exogenous sources.

Expression vectors generally have convenient restriction sites located near the promoter sequence to provide for the insertion of nucleic acid sequences encoding heterologous proteins. A selectable marker operative in the expression host may be present. Expression vectors may be used for the production of fusion proteins, where the exogenous fusion peptide provides additional functionality, i.e. increased protein synthesis, stability, reactivity with defined antisera, an enzyme marker, e.g. β -galactosidase, etc.

Expression cassettes may be prepared comprising a transcription initiation region, the gene or fragment thereof, and a transcriptional termination region. Of particular interest is the use of sequences that allow for the expression of functional epitopes or domains, usually at least about 8 amino acids in length, more usually at least about 15 amino acids in length, to about 25 amino acids, and up to the complete open reading frame of the gene. After introduction of the DNA, the cells containing the construct may be selected by means of a selectable marker, the cells expanded and then used for expression.

Subject proteins and polypeptides may be expressed in prokaryotes or eukaryotes in accordance with conventional ways, depending upon the purpose for expression. For large scale production of the protein, a unicellular organism, such as *E. coli*, *B. subtilis*, *S. cerevisiae*, insect cells in combination with baculovirus vectors, or cells of a higher organism such as vertebrates, particularly mammals, e.g. COS 7 cells, may be used as the expression host cells. In some situations, it is desirable to express the subject coding sequence in eukaryotic cells, where the DGAT2 α or MGAT1 protein will benefit from native folding and post-translational modifications. Small peptides can also be synthesized in the laboratory. Polypeptides that are subsets of the complete DGAT2 α or MGAT1 sequence may be used to identify and investigate parts of the protein important for function.

Once the source of the protein is identified and/or prepared, e.g. a transfected host expressing the protein is prepared, the protein is then purified to produce the desired DGAT2 α - or MGAT1- comprising composition. Any convenient protein purification procedures may be employed, where suitable protein purification methodologies are described in Guide to Protein Purification, (Deuthser ed.) (Academic Press, 1990). For

example, a lysate may prepared from the original source, e.g. naturally occurring cells or tissues that express DGAT2 α or the expression host expressing DGAT2 α , and purified using HPLC, exclusion chromatography, gel electrophoresis, affinity chromatography, and the like.

5 Specific expression systems of interest include bacterial, yeast, insect cell and mammalian cell derived expression systems. Representative systems from each of these categories is are provided below:

Bacteria. Expression systems in bacteria include those described in Chang *et al.*, *Nature* (1978) 275:615; Goeddel *et al.*, *Nature* (1979) 281:544; Goeddel *et al.*, *Nucleic Acids Res.* (1980) 8:4057; EP 0 036,776; U.S. Patent No. 4,551,433; DeBoer *et al.*, *Proc. Natl. Acad. Sci. (USA)* (1983) 80:21-25; and Siebenlist *et al.*, *Cell* (1980) 20:269.

Yeast. Expression systems in yeast include those described in Hinnen *et al.*, *Proc. Natl. Acad. Sci. (USA)* (1978) 75:1929; Ito *et al.*, *J. Bacteriol.* (1983) 153:163; Kurtz *et al.*, *Mol. Cell. Biol.* (1986) 6:142; Kunze *et al.*, *J. Basic Microbiol.* (1985) 25:141; Gleeson *et al.*, *J. Gen. Microbiol.* (1986) 132:3459; Roggenkamp *et al.*, *Mol. Gen. Genet.* (1986) 202:302; Das *et al.*, *J. Bacteriol.* (1984) 158:1165; De Louvencourt *et al.*, *J. Bacteriol.* (1983) 154:737; Van den Berg *et al.*, *Bio/Technology* (1990) 8:135; Kunze *et al.*, *J. Basic Microbiol.* (1985) 25:141; Cregg *et al.*, *Mol. Cell. Biol.* (1985) 5:3376; U.S. Patent Nos. 4,837,148 and 4,929,555; Beach and Nurse, *Nature* (1981) 300:706; Davidow *et al.*, *Curr. Genet.* (1985) 10:380; Gaillardin *et al.*, *Curr. Genet.* (1985) 10:49; Ballance *et al.*, *Biochem. Biophys. Res. Commun.* (1983) 112:284-289; Tilburn *et al.*, *Gene* (1983) 26:205-221; Yelton *et al.*, *Proc. Natl. Acad. Sci. (USA)* (1984) 81:1470-1474; Kelly and Hynes, *EMBO J.* (1985) 4:475479; EP 0 244,234; and WO 91/00357.

25 Insect Cells. Expression of heterologous genes in insects is accomplished as described in U.S. Patent No. 4,745,051; Friesen *et al.*, "The Regulation of Baculovirus Gene Expression", in: *The Molecular Biology Of Baculoviruses* (1986) (W. Doerfler, ed.); EP 0 127,839; EP 0 155,476; and Vlak *et al.*, *J. Gen. Virol.* (1988) 69:765-776; Miller *et al.*, *Ann. Rev. Microbiol.* (1988) 42:177; Carbonell *et al.*, *Gene* (1988) 73:409; Maeda *et al.*, *Nature* (1985) 315:592-594; Lebacqz-Verheyden *et al.*, *Mol. Cell. Biol.* (1988) 8:3129; Smith *et al.*, *Proc. Natl. Acad. Sci. (USA)* (1985) 82:8844; Miyajima *et al.*, *Gene* (1987) 58:273; and Martin *et al.*, *DNA* (1988) 7:99. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts are described in Luckow *et al.*,

Bio/Technology (1988) 6:47-55, Miller *et al.*, *Generic Engineering* (1986) 8:277-279, and Maeda *et al.*, *Nature* (1985) 315:592-594.

Mammalian Cells. Mammalian expression is accomplished as described in Dijkema *et al.*, *EMBO J.* (1985) 4:761, Gorman *et al.*, *Proc. Natl. Acad. Sci. (USA)* (1982) 79:6777, Boshart *et al.*, *Cell* (1985) 41:521 and U.S. Patent No. 4,399,216. Other features of mammalian expression are facilitated as described in Ham and Wallace, *Meth. Enz.* (1979) 58:44, Barnes and Sato, *Anal. Biochem.* (1980) 102:255, U.S. Patent Nos. 4,767,704, 4,657,866, 4,927,762, 4,560,655, WO 90/103430, WO 87/00195, and U.S. RE 30,985.

When any of the above host cells, or other appropriate host cells or organisms, are used to replicate and/or express the polynucleotides or nucleic acids of the invention, the resulting replicated nucleic acid, RNA, expressed protein or polypeptide, is within the scope of the invention as a product of the host cell or organism. The product is recovered by any appropriate means known in the art.

Once the gene corresponding to a selected polynucleotide is identified, its expression can be regulated in the cell to which the gene is native. For example, an endogenous gene of a cell can be regulated by an exogenous regulatory sequence as disclosed in U.S. Patent No. 5,641,670.

METHODS AND COMPOSITIONS HAVING RESEARCH APPLICATION

Also provided by the subject invention are methods and compositions having research applications, such as in the study of the acylglycerol metabolism; in the identification of key components of the di- and triglyceride synthesis pathway; in the identification of di- and triglyceride synthesis modulatory agents, e.g. DGAT2 α or MGAT1 inhibitors or enhancers, and the like.

The subject nucleic acid compositions find use in a variety of research applications. Research applications of interest include: the identification of DGAT2 α and MGAT1 homologs; as a source of novel promoter elements; the identification of DGAT2 α or MGAT1 expression regulatory factors; as probes and primers in hybridization applications, e.g. PCR; the identification of expression patterns in biological specimens; the preparation of cell or animal models for DGAT2 α or MGAT1 function; the preparation of *in vitro* models for DGAT2 α or MGAT1 function; etc.

Homologs of the specifically disclosed subject nucleic acids are identified by any of a number of methods. A fragment of the provided cDNA may be used as a hybridization probe against a cDNA library from the target organism of interest, where low stringency conditions are used. The probe may be a large fragment, or one or more short degenerate primers. Nucleic acids having sequence similarity are detected by hybridization under low stringency conditions, for example, at 50°C and 6×SSC (0.9 M sodium chloride/0.09 M sodium citrate) and remain bound when subjected to washing at 55°C in 1×SSC (0.15 M sodium chloride/0.015 M sodium citrate). Sequence identity may be determined by hybridization under stringent conditions, for example, at 50°C or higher and 0.1×SSC (15 mM sodium chloride/0.15 mM sodium citrate). Nucleic acids having a region of substantial identity to the provided nucleic acid sequences bind to the provided sequences under stringent hybridization conditions. By using probes, particularly labeled probes of DNA sequences, one can isolate homologous or related genes. One can also use sequence information derived from the polynucleotide compositions of the subject invention to prepare electronic “probes” for use in searching of computer based sequence data, e.g. BLAST searches EST databases.

The sequence of the 5' flanking region of the subject nucleic acid compositions may be utilized as a source for promoter elements, including enhancer-binding sites, that provide for developmental regulation in tissues where a subject acyltransferase, e.g., DGAT2 α or MGAT1, is expressed. The tissue-specific expression is useful for determining the pattern of expression, and for providing promoters that mimic the native pattern of expression. Naturally occurring polymorphisms in the promoter region are useful for determining natural variations in expression, particularly those that may be associated with disease.

Alternatively, mutations may be introduced into the promoter region to determine the effect of altering expression in experimentally defined systems. Methods for the identification of specific DNA motifs involved in the binding of transcriptional factors are known in the art, e.g. sequence similarity to known binding motifs, gel retardation studies, etc. For examples, see Blackwell *et al.* (1995), *Mol. Med.* 1:194-205; Mortlock *et al.* (1996), *Genome Res.* 6:327-33; and Joulin and Richard-Foy (1995), *Eur. J. Biochem.* 232:620-626.

The regulatory sequences may be used to identify *cis* acting sequences required for transcriptional or translational regulation of DGAT2 gene expression, especially in different tissues or stages of development, and to identify *cis* acting sequences and *trans*-acting factors that regulate or mediate DGAT2 or MGAT1 gene expression. Such transcription or

translational control regions may be operably linked to a DGAT2 or MGAT gene in order to promote expression of wild type or altered DGAT2 or MGAT1 or other proteins of interest in cultured cells, or in embryonic, fetal or adult tissues, and for gene therapy.

Small DNA fragments are useful as primers for PCR, hybridization screening probes,
5 *etc.* Larger DNA fragments, *i.e.* greater than 100 nucleotides (nt) are useful for production of the encoded polypeptide, as described in the previous section. For use in amplification reactions, such as PCR, a pair of primers will be used. The exact composition of the primer sequences is not critical to the invention, but for most applications the primers will hybridize to the subject sequence under stringent conditions, as known in the art. It is preferable to
10 choose a pair of primers that will generate an amplification product of at least about 50 nt, preferably at least about 100 nt. Algorithms for the selection of primer sequences are generally known, and are available in commercial software packages. Amplification primers hybridize to complementary strands of DNA, and will prime towards each other.

The DNA may also be used to identify expression of the gene in a biological
15 specimen. The manner in which one probes cells for the presence of particular nucleotide sequences, as genomic DNA or RNA, is well established in the literature. Briefly, DNA or mRNA is isolated from a cell sample. The mRNA may be amplified by RT-PCR, using reverse transcriptase to form a complementary DNA strand, followed by polymerase chain reaction amplification using primers specific for the subject DNA sequences. Alternatively,
20 the mRNA sample is separated by gel electrophoresis, transferred to a suitable support, *e.g.* nitrocellulose, nylon, *etc.*, and then probed with a fragment of the subject DNA as a probe. Other techniques, such as oligonucleotide ligation assays, *in situ* hybridizations, and hybridization to DNA probes arrayed on a solid chip may also find use. Detection of mRNA hybridizing to the subject sequence is indicative of DGAT2 α or MGAT1 gene expression in
25 the sample.

The sequence of a subject gene or nucleic acid, including flanking promoter regions and coding regions, may be mutated in various ways known in the art to generate targeted changes in promoter strength, sequence of the encoded protein, *etc.* The DNA sequence or protein product of such a mutation will usually be substantially similar to the sequences
30 provided herein, *i.e.* will differ by at least one nucleotide or amino acid, respectively, and may differ by at least two but not more than about ten nucleotides or amino acids. The sequence changes may be substitutions, insertions, deletions, or a combination thereof. Deletions may further include larger changes, such as deletions of a domain or exon. Other

modifications of interest include epitope tagging, e.g. with the FLAG system, HA, etc. For studies of subcellular localization, fusion proteins with green fluorescent proteins (GFP) may be used.

Techniques for *in vitro* mutagenesis of cloned genes are known. Examples of protocols for site specific mutagenesis may be found in Gustin *et al.* (1993), *Biotechniques* 14:22; Barany (1985), *Gene* 37:111-23; Colicelli *et al.* (1985), *Mol. Gen. Genet.* 199:537-9; and Prentki *et al.* (1984), *Gene* 29:303-13. Methods for site specific mutagenesis can be found in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, CSH Press 1989, pp. 15.3-15.108; Weiner *et al.* (1993), *Gene* 126:35-41; Sayers *et al.* (1992), *Biotechniques* 13:592-6; Jones and Winistorfer (1992), *Biotechniques* 12:528-30; Barton *et al.* (1990), *Nucleic Acids Res* 18:7349-55; Marotti and Tomich (1989), *Gene Anal. Tech.* 6:67-70; and Zhu (1989), *Anal Biochem* 177:120-4. Such mutated genes may be used to study structure-function relationships of DGAT2 α , or to alter properties of the protein that affect its function or regulation.

The subject nucleic acids can be used to generate transgenic hosts, e.g. non-human animals, such as mice, cows, rats, pigs etc., or site specific gene modifications in cell lines. Examples of transgenic hosts include hosts in which the naturally expressed DGAT2 α or MGAT1 gene has been disrupted, e.g. DGAT2 α or MGAT1 knock-outs, as well as hosts in which DGAT2 α or MGAT1 expression has been amplified, e.g. through introduction of additional DGAT2 α or MGAT1 copies, through introduction of strong promoter upstream of the DGAT2 α or MGAT1 gene, and the like. Using the nucleic acid compositions of the subject invention, standard protocols known to those of skill in the art may be used to produce such transgenic hosts that have been genetically manipulated with respect to the subject gene, i.e. DGAT2 α or MGAT1 transgenic hosts.

Transgenic animals may be made through homologous recombination, where the normal DGAT2 α or MGAT1 locus is altered, e.g. as in DGAT2 α or MGAT1 knockouts. Alternatively, a nucleic acid construct is randomly integrated into the genome. Vectors for stable integration include plasmids, retroviruses and other animal viruses, YACs, and the like. DNA constructs for homologous recombination will comprise at least a portion of the DGAT2 α or MGAT1 gene native to the species of the host animal, wherein the gene has the desired genetic modification(s), and includes regions of homology to the target locus. DNA constructs for random integration need not include regions of homology to mediate recombination. Conveniently, markers for positive and negative selection are included.

Methods for generating cells having targeted gene modifications through homologous recombination are known in the art. For various techniques for transfecting mammalian cells, see Keown *et al.* (1990), *Meth. Enzymol.* **185**:527-537.

For embryonic stem (ES) cells, an ES cell line may be employed, or embryonic cells
5 may be obtained freshly from a host, *e.g.* mouse, rat, guinea pig, cow, *etc.* Such cells are grown on an appropriate fibroblast-feeder layer or grown in the presence of leukemia inhibiting factor (LIF). When ES or embryonic cells have been transformed, they may be used to produce transgenic animals. After transformation, the cells are plated onto a feeder layer in an appropriate medium. Cells containing the construct may be detected by
10 employing a selective medium. After sufficient time for colonies to grow, they are picked and analyzed for the occurrence of homologous recombination or integration of the construct. Those colonies that are positive may then be used for embryo manipulation and blastocyst injection. Blastocysts are obtained from 4 to 6 week old superovulated females. The ES cells are trypsinized, and the modified cells are injected into the blastocoel of the
15 blastocyst. After injection, the blastocysts are returned to each uterine horn of pseudopregnant females. Females are then allowed to go to term and the resulting offspring screened for the construct. By providing for a different phenotype of the blastocyst and the genetically modified cells, chimeric progeny can be readily detected.

The resultant chimeric animals are screened for the presence of the modified gene
20 and males and females having the modification are mated to produce homozygous progeny. If the gene alterations cause lethality at some point in development, tissues or organs can be maintained as allogeneic or congenic grafts or transplants, or in *in vitro* culture. The transgenic animals may be any non-human mammal, such as laboratory animals, domestic animals, *etc.*

25 Transgenic plants may be produced in a similar manner. Methods of preparing transgenic plant cells and plants are described in U.S. Pat. Nos. 5,767,367; 5,750,870; 5,739,409; 5,689,049; 5,689,045; 5,674,731; 5,656,466; 5,633,155; 5,629,470 ; 5,595,896; 5,576,198; 5,538,879; 5,484,956; the disclosures of which are herein incorporated by reference. Methods of producing transgenic plants are also reviewed in Plant Biochemistry
30 and Molecular Biology (eds Lea & Leegood, John Wiley & Sons)(1993) pp 275-295. In brief, a suitable plant cell or tissue is harvested, depending on the nature of the plant species. As such, in certain instances, protoplasts will be isolated, where such protoplasts may be isolated from a variety of different plant tissues, *e.g.* leaf, hypocotyl, root, *etc.* For protoplast

isolation, the harvested cells are incubated in the presence of cellulases in order to remove the cell wall, where the exact incubation conditions vary depending on the type of plant and/or tissue from which the cell is derived. The resultant protoplasts are then separated from the resultant cellular debris by sieving and centrifugation. Instead of using protoplasts, embryogenic explants comprising somatic cells may be used for preparation of the transgenic host. Following cell or tissue harvesting, exogenous DNA of interest is introduced into the plant cells, where a variety of different techniques are available for such introduction. With isolated protoplasts, the opportunity arise for introduction via DNA-mediated gene transfer protocols, including: incubation of the protoplasts with naked DNA, e.g. plasmids, comprising the exogenous coding sequence of interest in the presence of polyvalent cations, e.g. PEG or PLO; and electroporation of the protoplasts in the presence of naked DNA comprising the exogenous sequence of interest. Protoplasts that have successfully taken up the exogenous DNA are then selected, grown into a callus, and ultimately into a transgenic plant through contact with the appropriate amounts and ratios of stimulatory factors, e.g. auxins and cytokinins. With embryogenic explants, a convenient method of introducing the exogenous DNA in the target somatic cells is through the use of particle acceleration or "gene-gun" protocols. The resultant explants are then allowed to grow into chimera plants, cross-bred and transgenic progeny are obtained. Instead of the naked DNA approaches described above, another convenient method of producing transgenic plants is *Agrobacterium* mediated transformation. With *Agrobacterium* mediated transformation, co-integrative or binary vectors comprising the exogenous DNA are prepared and then introduced into an appropriate *Agrobacterium* strain, e.g. *A. tumefaciens*. The resultant bacteria are then incubated with prepared protoplasts or tissue explants, e.g. leaf disks, and a callus is produced. The callus is then grown under selective conditions, selected and subjected to growth media to induce root and shoot growth to ultimately produce a transgenic plant.

The modified cells, animals or plants are useful in the study of function and regulation of a subject gene or nucleic acid. For example, a series of small deletions and/or substitutions may be made in the host's native DGAT2 α or MGAT1 gene to determine the role of different exons in various physiological processes. Specific constructs of interest include anti-sense nucleic acids, which will block DGAT2 α or MGAT1 expression, expression of dominant negative DGAT2 α or MGAT1 mutations, and over-expression of DGAT2 α or MGAT1 genes. Where a subject nucleic acid sequence is introduced, the

introduced sequence may be either a complete or partial sequence of a subject gene native to the host, or may be a complete or partial subject nucleic acid sequence that is exogenous to the host animal, *e.g.*, a human DGAT2 α or a human MGAT1 sequence. A detectable marker, such as *lac Z* (encoding β -galactosidase) may be introduced into the DGAT2 α or MGAT1 locus, where upregulation of DGAT2 α or MGAT1 gene expression will result in an easily detected change in phenotype. One may also provide for expression of the subject gene or variants thereof in cells or tissues where it is not normally expressed, at levels not normally present in such cells or tissues, or at abnormal times of development. The transgenic hosts, *e.g.* animals, plants, etc., may be used in functional studies, drug screening, *etc.*, *e.g.* to determine the effect of a candidate drug on DGAT2 α or MGAT1 activity, to identify drugs that reduce serum triglyceride levels, etc.

The subject polypeptide compositions can be used to produce *in vitro* models of diglyceride and/or triglyceride synthesis, where such models will consist of the subject proteins and other components of di- and/or triglyceride synthesis, *e.g.* substrates, such as monoacylglycerol, diacylglycerol or metabolic precursors thereof, fatty acyl CoAs and the like, other components of the diacylglycerol and/or triacylglycerol synthetase complex, *e.g.* acyl CoA ligase, acyl CoA acyltransferase, monoacyl glycerol acyltransferase, etc.

Also provided by the subject invention are screening assays designed to find modulatory agents of activity of a subject mono- or diacylglycerol acyltransferase, *e.g.* inhibitors or enhancers of DGAT2 α or MGAT1 activity, as well as the agents identified thereby, where such agents may find use in a variety of applications, including as therapeutic agents, as agricultural chemicals, etc. The screening methods will typically be assays which provide for qualitative/quantitative measurements of DGAT2 α or MGAT1 activity in the presence of a particular candidate therapeutic agent. For example, the assay could be an assay which measures the acylation activity of DGAT2 α or MGAT1 in the presence and absence of a candidate inhibitor agent. The screening method may be an *in vitro* or *in vivo* format, where both formats are readily developed by those of skill in the art.

Thus, in some embodiments, the invention provides an *in vitro* method of identifying an agent that modulates the acyltransferase activity of MGAT1. The method generally involves contacting MGAT1 with a candidate agent (also referred to as a "test agent") in the presence of an acyl donor and an acyl acceptor. The effect, if any, of a test agent on the amount of acylated acceptor that is produced is measured relative to a control sample, which control sample includes the MGAT1 polypeptide, the acyl donor, and the acyl acceptor, and

no test agent. Typically, the reaction mixture includes magnesium ions (e.g., $MgCl_2$); and a buffer. Exemplary reaction conditions are provided in Example 5. Suitable acyl donors are fatty acyl CoA compounds and include, but are not limited to, oleoyl CoA. Typically, the acyl group of the acyl donor is labeled with a detectable label, such that when the acyl group is transferred to the acyl acceptor, the detectable label is also transferred, thereby allowing detection of the acylated acceptor molecule. Suitable acyl acceptors are monoacylglycerols and diacylglycerols.

The DGAT2 α or MGAT1 polypeptide in the screening assay may be purified, but need not be. For example, membrane fractions containing DGAT2 α or MGAT1 polypeptides can be used.

Depending on the particular method, one or more of, usually one of, the components of the screening assay may be labeled, where by labeled is meant that the components comprise a detectable moiety, e.g. a fluorescent or radioactive tag, or a member of a signal producing system, e.g. biotin for binding to an enzyme-streptavidin conjugate in which the enzyme is capable of converting a substrate to a chromogenic product. Where in vitro assays are employed, the various components of the in vitro assay, e.g. the substrate, the donor, the DGAT2 α or MGAT1 protein and the candidate inhibitor, etc. are combined in a assay mixture under conditions sufficient for DGAT2 α or MGAT1 activity to occur, as described in the experimental section, *infra*.

A variety of other reagents may be included in the screening assay and reaction mixture. These include reagents like salts, neutral proteins, e.g. albumin, detergents, etc that are used to facilitate optimal protein-protein binding and/or reduce non-specific or background interactions. Reagents that improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, *etc.* may be used.

A variety of different candidate agents may be screened by the above methods. Candidate agents encompass numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 50 and less than about 2,500 daltons. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules

including peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof.

Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides and oligopeptides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means, and may be used to produce combinatorial libraries. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification, *etc.* to produce structural analogs.

Using the above screening methods, a variety of different therapeutic agents may be identified. Such agents may target the enzyme itself, or an expression regulator factor thereof. Such agents may be inhibitors or promoters of DGAT2 α or MGAT1 activity, where inhibitors are those agents that result in at least a reduction of DGAT2 α or MGAT1 activity as compared to a control and enhancers result in at least an increase in DGAT2 α or MGAT1 activity as compared to a control. Such agents may be found use in a variety of therapeutic applications, as described in greater detail below.

METHODS AND COMPOSITIONS HAVING MEDICAL APPLICATIONS

The methods and compositions of the subject invention also have broad ranging applications in a variety of medical applications, including diagnostic screening, therapeutic treatments of pathological conditions, in the regulation of DGAT2 α or MGAT1 activity in desirable ways, and the like.

The subject invention provides methods of screening individuals for a predisposition to a disease state or the presence of disease state, where such screening may focus on the presence of one or more markers, such as a mutated DGAT2 α or MGAT1 gene or expression regulatory element thereof, observed levels of DGAT2 α or MGAT1; the expression level of the DGAT2 α or MGAT1 gene in a biological sample of interest; and the like.

Samples, as used herein, include biological fluids such as blood, cerebrospinal fluid, tears, saliva, lymph, semen, dialysis fluid and the like; organ or tissue culture derived fluids; and fluids extracted from physiological tissues. Also included in the term are derivatives and fractions of such fluids. The cells may be dissociated, in the case of solid tissues, or tissue sections may be analyzed. Alternatively a lysate of the cells may be prepared.

A number of methods are available for determining the expression level of a gene or protein in a particular sample. Diagnosis may be performed by a number of methods to determine the absence or presence or altered amounts of normal or abnormal DGAT2 α or MGAT1 in a patient sample. For example, detection may utilize staining of cells or histological sections with labeled antibodies, performed in accordance with conventional methods. Cells are permeabilized to stain cytoplasmic molecules. The antibodies of interest are added to the cell sample, and incubated for a period of time sufficient to allow binding to the epitope, usually at least about 10 minutes. The antibody may be labeled with radioisotopes, enzymes, fluorescers, chemilumescers, or other labels for direct detection. Alternatively, a second stage antibody or reagent is used to amplify the signal. Such reagents are well known in the art. For example, the primary antibody may be conjugated to biotin, with horseradish peroxidase-conjugated avidin added as a second stage reagent. Alternatively, the secondary antibody conjugated to a fluorescent compound, *e.g.* fluorescein, rhodamine, Texas red, *etc.* Final detection uses a substrate that undergoes a color change in the presence of the peroxidase. The absence or presence of antibody binding may be determined by various methods, including flow cytometry of dissociated cells, microscopy, radiography, scintillation counting, *etc.*

Alternatively, one may focus on the expression of DGAT2 α - or MGAT1- encoding nucleic acids. Biochemical studies may be performed to determine whether a sequence polymorphism in a DGAT2 α or MGAT1 coding region or control regions is associated with disease. Disease associated polymorphisms may include deletion or truncation of the gene, mutations that alter expression level, that affect the activity of the protein, *etc.*

Changes in the promoter or enhancer sequence that may affect expression levels of DGAT2 α or MGAT1 can be compared to expression levels of the normal allele by various methods known in the art. Methods for determining promoter or enhancer strength include quantitation of the expressed natural protein; insertion of the variant control element into a vector with a reporter gene such as β -galactosidase, luciferase, chloramphenicol acetyltransferase, *etc.* that provides for convenient quantitation; and the like.

A number of methods are available for analyzing nucleic acids for the presence of a specific sequence, *e.g.* a disease associated polymorphism. Where large amounts of DNA are available, genomic DNA is used directly. Alternatively, the region of interest is cloned into a suitable vector and grown in sufficient quantity for analysis. Cells that express
5 DGAT2 α or MGAT1 may be used as a source of mRNA, which may be assayed directly or reverse transcribed into cDNA for analysis. The nucleic acid may be amplified by conventional techniques, such as the polymerase chain reaction (PCR), to provide sufficient amounts for analysis. The use of the polymerase chain reaction is described in Saiki, *et al.* (1985), *Science* 239:487, and a review of techniques may be found in Sambrook, *et al.*
10 Molecular Cloning: A Laboratory Manual, CSH Press 1989, pp.14.2-14.33. Alternatively, various methods are known in the art that utilize oligonucleotide ligation as a means of detecting polymorphisms, for examples see Riley *et al.* (1990), *Nucl. Acids Res.* 18:2887-2890; and Delahunty *et al.* (1996), *Am. J. Hum. Genet.* 58:1239-1246.

A detectable label may be included in an amplification reaction. Suitable labels
15 include fluorochromes, *e.g.* fluorescein isothiocyanate (FITC), rhodamine, Texas Red, phycoerythrin, allophycocyanin, 6-carboxyfluorescein (6-FAM), 2',7'-dimethoxy-4',5'-dichloro-6-carboxyfluorescein (JOE), 6-carboxy-X-rhodamine (ROX), 6-carboxy-2',4',7',4,7-hexachlorofluorescein (HEX), 5-carboxyfluorescein (5-FAM) or N,N,N',N'-tetramethyl-6-carboxyrhodamine (TAMRA), radioactive labels, *e.g.* ^{32}P , ^{35}S , ^3H ; *etc.* The
20 label may be a two stage system, where the amplified DNA is conjugated to biotin, haptens, *etc.* having a high affinity binding partner, *e.g.* avidin, specific antibodies, *etc.*, where the binding partner is conjugated to a detectable label. The label may be conjugated to one or both of the primers. Alternatively, the pool of nucleotides used in the amplification is labeled, so as to incorporate the label into the amplification product.

25 The sample nucleic acid, *e.g.* amplified or cloned fragment, is analyzed by one of a number of methods known in the art. The nucleic acid may be sequenced by dideoxy or other methods, and the sequence of bases compared to a wild-type DGAT2 α or MGAT1 sequence. Hybridization with the variant sequence may also be used to determine its presence, by Southern blots, dot blots, *etc.* The hybridization pattern of a control and variant
30 sequence to an array of oligonucleotide probes immobilized on a solid support, as described in US 5,445,934, or in WO 95/35505, may also be used as a means of detecting the presence of variant sequences. Single strand conformational polymorphism (SSCP) analysis, denaturing gradient gel electrophoresis (DGGE), and heteroduplex analysis in gel matrices

are used to detect conformational changes created by DNA sequence variation as alterations in electrophoretic mobility. Alternatively, where a polymorphism creates or destroys a recognition site for a restriction endonuclease, the sample is digested with that endonuclease, and the products size fractionated to determine whether the fragment was digested.

- 5 Fractionation is performed by gel or capillary electrophoresis, particularly acrylamide or agarose gels.

Screening for mutations in DGAT2 α or MGAT1 may be based on the functional or antigenic characteristics of the protein. Protein truncation assays are useful in detecting deletions that may affect the biological activity of the protein. Various immunoassays
10 designed to detect polymorphisms in DGAT2 α or MGAT1 proteins may be used in screening. Where many diverse genetic mutations lead to a particular disease phenotype, functional protein assays have proven to be effective screening tools. The activity of the encoded DGAT2 α or MGAT1 protein may be determined by comparison with the wild-type protein.

15 Diagnostic methods of the subject invention in which the level of DGAT2 α or MGAT1 expression is of interest will typically involve comparison of the DGAT2 α or MGAT1 nucleic acid abundance of a sample of interest with that of a control value to determine any relative differences, where the difference may be measured qualitatively and/or quantitatively, which differences are then related to the presence or absence of an
20 abnormal DGAT2 α or MGAT1 expression pattern. A variety of different methods for determining the nucleic acid abundance in a sample are known to those of skill in the art, where particular methods of interest include those described in: Pietu et al., Genome Res. (June 1996) 6: 492-503; Zhao et al., Gene (April 24, 1995) 156: 207-213; Soares, Curr. Opin. Biotechnol. (October 1997) 8: 542-546; Raval, J. Pharmacol Toxicol Methods
25 (November 1994) 32: 125-127; Chalifour et al., Anal. Biochem (February 1, 1994) 216: 299-304; Stolz & Tuan, Mol. Biotechnol. (December 1996) 6: 225-230; Hong et al., Bioscience Reports (1982) 2: 907; and McGraw, Anal. Biochem. (1984) 143: 298. Also of interest are the methods disclosed in WO 97/27317, the disclosure of which is herein incorporated by reference.

30 The subject diagnostic or screening methods may be used to identify the presence of, or predisposition to, disease conditions associated with acylglycerol metabolism, particularly those associated with DGAT2 α or MGAT1 activity. Such disease conditions include: hyperlipidemia (including excess serum triglyceride levels), cardiovascular disease, obesity,

diabetes, cancer, neurological disorders, immunological disorders, and the like.

Also provided are methods of regulating, including enhancing and inhibiting, DGAT2 α or MGAT1 activity in a host. A variety of situations arise where modulation of DGAT2 α or MGAT1 activity in a host is desired, where such conditions include disease
5 conditions associated with DGAT2 α or MGAT1 activity and non-disease conditions in which a modulation of DGAT2 α or MGAT1 activity is desired for a variety of different reasons, e.g. cosmetic weight control.

For the modulation of DGAT2 α or MGAT1 activity in a host, an effective amount of active agent that modulates the activity, e.g. reduces the activity, of DGAT2 α or MGAT1 *in*
10 *vivo*, is administered to the host. The active agent may be a variety of different compounds, including: the polynucleotide compositions of the subject invention, the polypeptide compositions of the subject invention, a naturally occurring or synthetic small molecule compound, an antibody, fragment or derivative thereof, an antisense composition, and the like.

15 The nucleic acid compositions of the subject invention find use as therapeutic agents in situations where one wishes to enhance DGAT2 α or MGAT1 activity in a host, e.g. in a mammalian host in which DGAT2 α or MGAT1 activity is low resulting in a disease condition, etc. The DGAT2 α or MGAT1 genes, gene fragments, or the encoded DGAT2 α or MGAT1 protein or protein fragments are useful in gene therapy to treat disorders associated
20 with DGAT2 α or MGAT1 defects. Expression vectors may be used to introduce the DGAT2 α gene or encoding nucleic acid into a cell. Such vectors generally have convenient restriction sites located near the promoter sequence to provide for the insertion of nucleic acid sequences. Transcription cassettes may be prepared comprising a transcription initiation region, the target gene or fragment thereof, and a transcriptional termination
25 region. The transcription cassettes may be introduced into a variety of vectors, e.g. plasmid; retrovirus, e.g. lentivirus; adenovirus; and the like, where the vectors are able to transiently or stably be maintained in the cells, usually for a period of at least about one day, more usually for a period of at least about several days to several weeks.

Naturally occurring or synthetic small molecule compounds of interest include
30 numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 50 and less than about 2,500 daltons. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding, and typically include at least an amine,

carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules including peptides, 5 saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof. Of particular interest are those agents identified by the screening assays of the subject invention, as described above.

Also of interest as active agents are antibodies that modulate, e.g. reduce, if not inhibit, DGAT2 α or MGAT1 activity in the host. Suitable antibodies are obtained by 10 immunizing a host animal with peptides comprising all or a portion of a DGAT2 or MGAT1 protein, such as the DGAT2 α or MGAT1 polypeptide compositions of the subject invention. Suitable host animals include mouse, rat sheep, goat, hamster, rabbit, *etc.* The origin of the protein immunogen may be mouse, human, rat, monkey *etc.* The host animal will generally be a different species than the immunogen, e.g. human DGAT2 used to immunize mice, *etc.*

15 The immunogen may comprise the complete protein, or fragments and derivatives thereof. Preferred immunogens comprise all or a part of DGAT2 α or MGAT1, where these residues contain the post-translation modifications, such as glycosylation, found on the native DGAT2 α or MGAT1. Immunogens comprising the extracellular domain are produced in a variety of ways known in the art, e.g. expression of cloned genes using conventional 20 recombinant methods, isolation from HEC, *etc.*

For preparation of polyclonal antibodies, the first step is immunization of the host animal with DGAT2 α or MGAT1, where the DGAT2 α or MGAT1 protein will preferably be in substantially pure form, comprising less than about 1% contaminant. The immunogen may comprise complete DGAT2 α or MGAT1, fragments or derivatives thereof. To increase 25 the immune response of the host animal, the DGAT2 or MGAT1 may be combined with an adjuvant, where suitable adjuvants include alum, dextran, sulfate, large polymeric anions, oil & water emulsions, e.g. Freund's adjuvant, Freund's complete adjuvant, and the like. The DGAT2 α or MGAT1 may also be conjugated to synthetic carrier proteins or synthetic antigens. A variety of hosts may be immunized to produce the polyclonal antibodies. Such 30 hosts include rabbits, guinea pigs, rodents, e.g. mice, rats, sheep, goats, and the like. The DGAT2 α or MGAT1 is administered to the host, usually intradermally, with an initial dosage followed by one or more, usually at least two, additional booster dosages. Following immunization, the blood from the host will be collected, followed by separation of the serum

from the blood cells. The Ig present in the resultant antiserum may be further fractionated using known methods, such as ammonium salt fractionation, DEAE chromatography, and the like.

Monoclonal antibodies are produced by conventional techniques. Generally, the spleen and/or lymph nodes of an immunized host animal provide a source of plasma cells. The plasma cells are immortalized by fusion with myeloma cells to produce hybridoma cells. Culture supernatant from individual hybridomas is screened using standard techniques to identify those producing antibodies with the desired specificity. Suitable animals for production of monoclonal antibodies to the human protein include mouse, rat, hamster, *etc.* To raise antibodies against the mouse protein, the animal will generally be a hamster, guinea pig, rabbit, *etc.* The antibody may be purified from the hybridoma cell supernatants or ascites fluid by conventional techniques, *e.g.* affinity chromatography using DGAT2 α or MGAT1 bound to an insoluble support, protein A sepharose, *etc.*

The antibody may be produced as a single chain, instead of the normal multimeric structure. Single chain antibodies are described in Jost *et al.* (1994) J.B.C. 269:26267-73, and others. DNA sequences encoding the variable region of the heavy chain and the variable region of the light chain are ligated to a spacer encoding at least about 4 amino acids of small neutral amino acids, including glycine and/or serine. The protein encoded by this fusion allows assembly of a functional variable region that retains the specificity and affinity of the original antibody.

For *in vivo* use, particularly for injection into humans, it is desirable to decrease the antigenicity of the antibody. An immune response of a recipient against the blocking agent will potentially decrease the period of time that the therapy is effective. Methods of humanizing antibodies are known in the art. The humanized antibody may be the product of an animal having transgenic human immunoglobulin constant region genes (see for example International Patent Applications WO 90/10077 and WO 90/04036). Alternatively, the antibody of interest may be engineered by recombinant DNA techniques to substitute the CH1, CH2, CH3, hinge domains, and/or the framework domain with the corresponding human sequence (see WO 92/02190).

The use of Ig cDNA for construction of chimeric immunoglobulin genes is known in the art (Liu *et al.* (1987) P.N.A.S. 84:3439 and (1987) J. Immunol. 139:3521). mRNA is isolated from a hybridoma or other cell producing the antibody and used to produce cDNA. The cDNA of interest may be amplified by the polymerase chain reaction using specific

primers (U.S. Patent nos. 4,683,195 and 4,683,202). Alternatively, a library is made and screened to isolate the sequence of interest. The DNA sequence encoding the variable region of the antibody is then fused to human constant region sequences. The sequences of human constant regions genes may be found in Kabat *et al.* (1991) Sequences of Proteins of Immunological Interest, N.I.H. publication no. 91-3242. Human C region genes are readily available from known clones. The choice of isotype will be guided by the desired effector functions, such as complement fixation, or activity in antibody-dependent cellular cytotoxicity. Preferred isotypes are IgG1, IgG3 and IgG4. Either of the human light chain constant regions, kappa or lambda, may be used. The chimeric, humanized antibody is then expressed by conventional methods.

Antibody fragments, such as Fv, F(ab')₂ and Fab may be prepared by cleavage of the intact protein, *e.g.* by protease or chemical cleavage. Alternatively, a truncated gene is designed. For example, a chimeric gene encoding a portion of the F(ab')₂ fragment would include DNA sequences encoding the CH1 domain and hinge region of the H chain, followed by a translational stop codon to yield the truncated molecule.

Consensus sequences of H and L J regions may be used to design oligonucleotides for use as primers to introduce useful restriction sites into the J region for subsequent linkage of V region segments to human C region segments. C region cDNA can be modified by site directed mutagenesis to place a restriction site at the analogous position in the human sequence.

Expression vectors include plasmids, retroviruses, YACs, EBV derived episomes, and the like. A convenient vector is one that encodes a functionally complete human CH or CL immunoglobulin sequence, with appropriate restriction sites engineered so that any VH or VL sequence can be easily inserted and expressed. In such vectors, splicing usually occurs between the splice donor site in the inserted J region and the splice acceptor site preceding the human C region, and also at the splice regions that occur within the human CH exons. Polyadenylation and transcription termination occur at native chromosomal sites downstream of the coding regions. The resulting chimeric antibody may be joined to any strong promoter, including retroviral LTRs, *e.g.* SV-40 early promoter, (Okayama *et al.* (1983) Mol. Cell. Bio. 3:280), Rous sarcoma virus LTR (Gorman *et al.* (1982) P.N.A.S. 79:6777), and moloney murine leukemia virus LTR (Grosschedl *et al.* (1985) Cell 41:885); native Ig promoters, *etc.*

In yet other embodiments of the invention, the active agent is an agent that modulates, and generally decreases or down regulates, the expression of DGAT2 α - or MGAT1-encoding nucleic acids in the host. Antisense molecules can be used to down-regulate expression of these target nucleic acids in cells. The anti-sense reagent may be
5 antisense oligonucleotides (ODN), particularly synthetic ODN having chemical modifications from native nucleic acids, or nucleic acid constructs that express such antisense molecules as RNA. The antisense sequence is complementary to the mRNA of the targeted gene, and inhibits expression of the targeted gene products. Antisense molecules inhibit gene expression through various mechanisms, *e.g.* by reducing the amount of mRNA
10 available for translation, through activation of RNase H, or steric hindrance. One or a combination of antisense molecules may be administered, where a combination may comprise multiple different sequences.

Antisense molecules may be produced by expression of all or a part of the target gene sequence in an appropriate vector, where the transcriptional initiation is oriented such that an
15 antisense strand is produced as an RNA molecule. Alternatively, the antisense molecule is a synthetic oligonucleotide. Antisense oligonucleotides will generally be at least about 7, usually at least about 12, more usually at least about 20 nucleotides in length, and not more than about 500, usually not more than about 50, more usually not more than about 35 nucleotides in length, where the length is governed by efficiency of inhibition, specificity,
20 including absence of cross-reactivity, and the like. It has been found that short oligonucleotides, of from 7 to 8 bases in length, can be strong and selective inhibitors of gene expression (see Wagner *et al.* (1996), *Nature Biotechnol.* 14:840-844).

A specific region or regions of the endogenous sense strand mRNA sequence is chosen to be complemented by the antisense sequence. Selection of a specific sequence for
25 the oligonucleotide may use an empirical method, where several candidate sequences are assayed for inhibition of expression of the target gene in an *in vitro* or animal model. A combination of sequences may also be used, where several regions of the mRNA sequence are selected for antisense complementation.

30 Antisense oligonucleotides may be chemically synthesized by methods known in the art (see Wagner *et al.* (1993), *supra*, and Milligan *et al.*, *supra*.) Preferred oligonucleotides are chemically modified from the native phosphodiester structure, in order to increase their intracellular stability and binding affinity. A number of such modifications have been

described in the literature, which alter the chemistry of the backbone, sugars or heterocyclic bases.

Among useful changes in the backbone chemistry are phosphorothioates; phosphorodithioates, where both of the non-bridging oxygens are substituted with sulfur; phosphoroamidites; alkyl phosphotriesters and boranophosphates. Achiral phosphate derivatives include 3'-O'-5'-S-phosphorothioate, 3'-S-5'-O-phosphorothioate, 3'-CH₂-5'-O-phosphonate and 3'-NH-5'-O-phosphoroamidate. Peptide nucleic acids replace the entire ribose phosphodiester backbone with a peptide linkage. Sugar modifications are also used to enhance stability and affinity. The α -anomer of deoxyribose may be used, where the base is inverted with respect to the natural β -anomer. The 2'-OH of the ribose sugar may be altered to form 2'-O-methyl or 2'-O-allyl sugars, which provides resistance to degradation without comprising affinity. Modification of the heterocyclic bases must maintain proper base pairing. Some useful substitutions include deoxyuridine for deoxythymidine; 5-methyl-2'-deoxycytidine and 5-bromo-2'-deoxycytidine for deoxycytidine. 5-propynyl-2'-deoxyuridine and 5-propynyl-2'-deoxycytidine have been shown to increase affinity and biological activity when substituted for deoxythymidine and deoxycytidine, respectively.

As an alternative to anti-sense inhibitors, catalytic nucleic acid compounds, *e.g.* ribozymes, anti-sense conjugates, *etc.* may be used to inhibit gene expression. Ribozymes may be synthesized *in vitro* and administered to the patient, or may be encoded on an expression vector, from which the ribozyme is synthesized in the targeted cell (for example, see International patent application WO 9523225, and Beigelman *et al.* (1995), *Nucl. Acids Res.* 23:4434-42). Examples of oligonucleotides with catalytic activity are described in WO 9506764. Conjugates of anti-sense ODN with a metal complex, *e.g.* terpyridylCu(II), capable of mediating mRNA hydrolysis are described in Bashkin *et al.* (1995), *Appl. Biochem. Biotechnol.* 54:43-56.

As mentioned above, an effective amount of the active agent is administered to the host, where "effective amount" means a dosage sufficient to produce a desired result, where the desired result in the desired modulation, *e.g.* enhancement, reduction, of DGAT2 α activity.

In the subject methods, the active agent(s) may be administered to the host using any convenient means capable of resulting in the desired effect. Thus, the agent can be incorporated into a variety of formulations for therapeutic administration. More particularly, the agents of the present invention can be formulated into pharmaceutical compositions by

combination with appropriate, pharmaceutically acceptable carriers or diluents, and may be formulated into preparations in solid, semi-solid, liquid or gaseous forms, such as tablets, capsules, powders, granules, ointments, solutions, suppositories, injections, inhalants and aerosols.

5 As such, administration of the agents can be achieved in various ways, including oral, buccal, rectal, parenteral, intraperitoneal, intradermal, transdermal, intracheal, etc., administration.

 In pharmaceutical dosage forms, the agents may be administered in the form of their pharmaceutically acceptable salts, or they may also be used alone or in appropriate
10 association, as well as in combination, with other pharmaceutically active compounds. The following methods and excipients are merely exemplary and are in no way limiting.

 For oral preparations, the agents can be used alone or in combination with appropriate additives to make tablets, powders, granules or capsules, for example, with conventional additives, such as lactose, mannitol, corn starch or potato starch; with binders,
15 such as crystalline cellulose, cellulose derivatives, acacia, corn starch or gelatins; with disintegrators, such as corn starch, potato starch or sodium carboxymethylcellulose; with lubricants, such as talc or magnesium stearate; and if desired, with diluents, buffering agents, moistening agents, preservatives and flavoring agents.

 The agents can be formulated into preparations for injection by dissolving,
20 suspending or emulsifying them in an aqueous or nonaqueous solvent, such as vegetable or other similar oils, synthetic aliphatic acid glycerides, esters of higher aliphatic acids or propylene glycol; and if desired, with conventional additives such as solubilizers, isotonic agents, suspending agents, emulsifying agents, stabilizers and preservatives.

 The agents can be utilized in aerosol formulations to be administered via inhalation.
25 The compounds of the present invention can be formulated into pressurized acceptable propellants such as dichlorodifluoromethane, propane, nitrogen and the like.

 Furthermore, the agents can be made into suppositories by mixing with a variety of bases such as emulsifying bases or water-soluble bases. The compounds of the present invention can be administered rectally via a suppository. The suppository can include
30 vehicles such as cocoa butter, carbowaxes and polyethylene glycols, which melt at body temperature, yet are solidified at room temperature.

 Unit dosage forms for oral or rectal administration such as syrups, elixirs, and suspensions may be provided wherein each dosage unit, for example, teaspoonful,

tablespoonful, tablet or suppository, contains a predetermined amount of the composition containing one or more inhibitors. Similarly, unit dosage forms for injection or intravenous administration may comprise the inhibitor(s) in a composition as a solution in sterile water, normal saline or another pharmaceutically acceptable carrier.

5 The term "unit dosage form," as used herein, refers to physically discrete units suitable as unitary dosages for human and animal subjects, each unit containing a predetermined quantity of compounds of the present invention calculated in an amount sufficient to produce the desired effect in association with a pharmaceutically acceptable diluent, carrier or vehicle. The specifications for the novel unit dosage forms of the present invention depend on the particular compound employed and the effect to be achieved, and
10 the pharmacodynamics associated with each compound in the host.

 The pharmaceutically acceptable excipients, such as vehicles, adjuvants, carriers or diluents, are readily available to the public. Moreover, pharmaceutically acceptable auxiliary substances, such as pH adjusting and buffering agents, tonicity adjusting agents, stabilizers,
15 wetting agents and the like, are readily available to the public.

 Where the agent is a polypeptide, polynucleotide, analog or mimetic thereof, e.g. antisense composition, it may be introduced into tissues or host cells by any number of routes, including viral infection, microinjection, or fusion of vesicles. Jet injection may also be used for intramuscular administration, as described by Furth *et al.* (1992), *Anal Biochem*
20 205:365-368. The DNA may be coated onto gold microparticles, and delivered intradermally by a particle bombardment device, or "gene gun" as described in the literature (see, for example, Tang *et al.* (1992), *Nature* 356:152-154), where gold microprojectiles are coated with the *DGAT* DNA, then bombarded into skin cells.

 Those of skill will readily appreciate that dose levels can vary as a function of the
25 specific compound, the severity of the symptoms and the susceptibility of the subject to side effects. Preferred dosages for a given compound are readily determinable by those of skill in the art by a variety of means.

 The subject methods find use in the treatment of a variety of different disease
30 conditions involving acylglycerol metabolism, and particularly DGAT2 α activity, including both insufficient or hypo- DGAT2 α activity and hyper- DGAT2 α activity. Representative diseases that may be treated according to the subject methods include: hyperlipidemia (including excess serum triglyceride levels), cardiovascular disease, obesity, diabetes,

cancer, neurological disorders, immunological disorders, skin disorders associated with sebaceous gland activity, e.g. acne, and the like.

By treatment is meant at least an amelioration of the symptoms associated with the pathological condition afflicting the host, where amelioration is used in a broad sense to refer to at least a reduction in the magnitude of a parameter, e.g. symptom, associated with the pathological condition being treated, such as serum triglyceride level, weight, total body fat content, etc. As such, treatment also includes situations where the pathological condition, or at least symptoms associated therewith, are completely inhibited, e.g. prevented from happening, or stopped, e.g. terminated, such that the host no longer suffers from the pathological condition, or at least the symptoms that characterize the pathological condition. For example, where the disease condition is marked by the presence of elevated lipid levels, treatment includes at least a reduction in the observed lipid levels, including a restoration of normal lipid levels. As another example, where the disease is obesity, treatment results in at least a reduction in the overall weight and/or total body fat content of the host.

The subject methods also find use in the modulation of DGAT2 α or MGAT1 activity in hosts not suffering from a disease condition but in which the modulation of DGAT2 α or MGAT1 activity is nonetheless desired. For example, sperm production in males has been associated with diglyceride acyltransferase activity. As such, in males where at least reduced sperm production is desired, the subject methods can be used to reduce this target activity in such males, e.g. by administering an agent that reduces DGAT2 α activity in such males, where such agents are described above. In other words, the subject methods provide a means of male contraception. Alternatively, where increased sperm count in a given male is desired, e.g. in those conditions where the male has reduced fertility, the subject methods can be used to enhance this target activity in the male and thereby increase sperm count and fertility, e.g. by administering to the male host a DGAT2 α enhancing agent, as described above.

A variety of hosts are treatable according to the subject methods. Generally such hosts are "mammals" or "mammalian," where these terms are used broadly to describe organisms which are within the class mammalia, including the orders carnivore (e.g., dogs and cats), rodentia (e.g., mice, guinea pigs, and rats), and primates (e.g., humans, chimpanzees, and monkeys). In many embodiments, the hosts will be humans.

Kits with unit doses of the active agent, usually in oral or injectable doses, are provided. In such kits, in addition to the containers containing the unit doses will be an informational package insert describing the use and attendant benefits of the drugs in treating

pathological condition of interest. Preferred compounds and unit doses are those described herein above.

METHODS AND COMPOSITIONS FOR PRODUCING DIGLYCERIDES, DIGLYCERIDE COMPOSITIONS,
5 TRIGLYCERIDES, AND TRIGLYCERIDE COMPOSITIONS

Also provided by the subject invention are methods for preparing diglycerides, diglyceride compositions, triglycerides, and triglyceride comprising compositions, as well as the compositions produced by these methods.

10 In preparing triglycerides with the subject invention, at least the direct substrates of the desired triacylglycerol, e.g. diacylglycerol and fatty acyl CoA, will be combined in the presence of the polypeptide under conditions sufficient for the acylation of the diacylglycerol to occur. The synthesis may occur in an *in vitro* system, e.g. in a vessel in which the substrates or precursors thereof and the DGAT2 α enzyme, as well as any other requisite enzymes (e.g. as need to convert the substrate precursors to substrates), or an *in vivo* system,
15 e.g. a host cell that naturally comprises the substrates and into which a DGAT2 α gene or nucleic acid according to the subject invention has been inserted in a manner sufficient for expression of the gene and provision of the DGAT2 α enzyme, where the resultant triglyceride products may be separated from the host cell using standard separation techniques.

20 In preparing diglycerides with the subject invention, at least the direct substrates of the desired diacylglycerol, e.g. monoacylglycerol and fatty acyl CoA, will be combined in the presence of the polypeptide under conditions sufficient for the acylation of the monoacylglycerol to occur. The synthesis may occur in an *in vitro* system, e.g. in a vessel in which the substrates or precursors thereof and the MGAT1 enzyme, as well as any other
25 requisite enzymes (e.g. as need to convert the substrate precursors to substrates), or an *in vivo* system, e.g. a host cell that naturally comprises the substrates and into which an MGAT1 gene or nucleic acid according to the subject invention has been inserted in a manner sufficient for expression of the gene and provision of the MGAT1 enzyme, where the resultant diglyceride products may be separated from the host cell using standard
30 separation techniques.

Of interest for use in producing di- and triglyceride compositions are transgenic plants/fungi that have been genetically manipulated using the nucleic acid compositions of the subject invention to produce di- and/or triglycerides and/or compositions thereof in one

or more desirable ways. Transgenic plants/fungi of the subject invention are those plants/fungi that at least: (a) produce more diglyceride, diglyceride composition, triglyceride or triglyceride composition than wild type, e.g. produce more oil, such as by producing seeds having a higher oil content, as compared to wild-type; (b) produce di- or triglyceride compositions, e.g. oils, that are enriched for di- or triglycerides and/or enriched for one or more particular di- or triglycerides as compared to wild type; and the like. Of interest are transgenic plants that produce commercially valuable triglyceride compositions or oils, such as canola, rapeseed, palm, corn, etc., containing various poly- and mono-unsaturated fatty acids, and the like. Of particular interest are transgenic plants, such as canola, rapeseed, palm, oil, etc., which have been genetically modified to produce seeds having higher oil content than the content found in the corresponding wild type, where the oil content of the seeds produced by such plants is at least 10% higher, usually at least 20 % higher, and in many embodiments at least 30 % higher than that found in the wild type, where in many embodiments seeds having oil contents that are 50% higher, or even greater, as compared to seeds produced by the corresponding wild-type plant, are produced. The seeds produced by such DGAT2 α transgenic plants can be used as sources of oil or as sources of additional DGAT2 α transgenic plants. Such transgenic plants and seeds therefore find use in methods of producing oils. In such methods, DGAT2 α transgenic plants engineered to produce seeds having a higher oil content than the corresponding wild-type, e.g. seeds in which the DGAT2 α gene is overexpressed, are grown, the seeds are harvested and then processed to recover the oil. The subject transgenic plants can also be used to produce novel oils characterized by the presence of triglycerides in different amounts and/or ratios than those observed in naturally occurring oils. The transgenic plants/fungi described above can be readily produced by those of skill in the art armed with the nucleic acid compositions of the subject invention. See the discussion on how to prepare transgenic plants, *supra*.

The triglyceride compositions described above find use in a variety of different applications. For example, such compositions or oils find use as food stuffs, being used as ingredients, spreads, cooking materials, etc. Alternatively, such oils find use as industrial feedstocks for use in the production of chemicals, lubricants, surfactants and the like.

Also of interest are transgenic non-human animals suitable for use as sources of food products and/or animal based industrial products. Such transgenic non-human animals, e.g. transgenic mice, rats, livestock, such as cows, pigs, horses, birds, etc, may be produced using methods known in the art and reviewed *supra*. Such transgenic non-human animals can be

used for sources of a variety of different food and industrial products in which the triglyceride content is specifically tailored in a desirable manner. For example, such transgenic animals that have been modified in a manner such that DGAT2 α activity is reduced as compared to the wild type can be used as sources of food products that are low in triglyceride content, e.g. low fat or lean meat products, low fat milk, low fat eggs, and the like.

The following examples are offered primarily for purposes of illustration. It will be readily apparent to those skilled in the art that the formulations, dosages, methods of administration, and other parameters of this invention may be further modified or substituted in various ways without departing from the spirit and scope of the invention.

EXPERIMENTAL

Example 1: Existence of DGAT2 α

A. Mice (DGAT1 $^{-/-}$) lacking DGAT, as described in WO 99/67268 are lean and resistant to diet-induced obesity, but are still capable of synthesizing triglycerides (TG) and have normal plasma TG levels. However, DGAT activity is virtually absent in membrane preparations from DGAT1 $^{-/-}$ tissues (Smith et al., Nat.Genet. 2000 (25), 87-90). Using pulse assays in living cells, we measured that the residual TG synthesis activity in DGAT1 $^{-/-}$ Mouse Embryonic Fibroblasts (MEF) or adipocytes was about 40% of that in wild-type cells. The results are graphically depicted in Figs. 1A and 1B. In Fig. 1A the membrane fraction isolated from MEF or adipocytes of wild-type or DGAT1 $^{-/-}$ mice was used as the enzyme source in DGAT assays *in vitro*. In Fig. 1B living cells were pulse-labeled with [14 C]oleic acid for 24 hours and [14 C] incorporation in the TG fraction was measured.

In further assays, increased DGAT activity was observed in DGAT1 $^{-/-}$ membranes assayed without magnesium; and DGAT activity was observed to vary with magnesium concentration in liver and adipose tissue.

The above findings indicate the existence of DGAT2 α , a second enzyme with diglyceride acyltransferase activity.

II. Mammalian DGAT2 α Sequences

A. The human DGAT2 α nucleic acid and amino acid sequences were identified using standard procedures, as described above. The human DGAT2 α cDNA has the sequence appearing as SEQ ID NO:01, infra, while the protein encoded thereby has the sequence appearing as SEQ ID NO:02, infra.

B. The mouse DGAT2 α nucleic acid and amino acid sequences were identified using standard procedures, as described above. The mouse DGAT2 α cDNA has the sequence appearing as SEQ ID NO:03, infra, while the protein encoded thereby has the sequence appearing as SEQ ID NO:04, infra.

Example 2: Characterization of DGAT2 α

A. Molecular Weight

The mouse DGAT2 α cDNA was determined to encode a 43 kD predicted protein based on the amino acid sequence. The mouse DGAT2 α cDNA was determined to have no sequence homology to DGAT1, as described in Cases et al., supra. The mouse DGAT2 α amino acid sequence was determined to have 2 putative N-linked glycosylation sites. The mouse DGAT2 α amino acid sequence was determined to have 6 putative PKC phosphorylation sites. A Hydrophobicity plot assessed by Kyte-Doolittle (K-D) analysis revealed the existence of multiple putative transmembrane domains in the mouse DGAT2 α amino acid sequence. Fig. 2 provides a graphical result of this analysis. As such, there are regions of higher hydrophobicity compatible with the existence of one or more transmembrane domain.

Example 3: Expression of DGAT2a in insect cells

Sf9 insect cells were infected with wild-type baculovirus, mouse FLAG-tagged DGAT2 α or mouse FLAG-tagged DGAT1 (Cases et al., supra) recombinant baculoviruses, and the membrane fractions were assayed for DGAT activity. The results are graphically provided in Fig. 3A. In Fig. 3A a time course of DGAT2 α virus infection is provided. Insect cell membranes were isolated at the indicated times after infection. Expression of the FLAG-

tagged DGAT2 α protein was detected by immunoblotting with an anti-FLAG antibody (*Inset*). DGAT activity was measured at low (5 mM) or high (100 mM) magnesium concentration, using [14 C]oleoyl CoA and cold diacylglycerol. The experiment was repeated three times and a representative experiment is shown. Fig. 3B shows that DGAT2 α activity is dependent on the presence of the diacylglycerol substrate. Assays were performed at low magnesium concentration, using [14 C]oleoyl CoA with or without exogenous cold diacylglycerol. When no diacylglycerol is added, no significant DGAT activity can be detected over background. Data represent the mean (\pm SD) of three experiments. To compare the DGAT activity of DGAT1 and DGAT2 α , membranes expressing equal levels of DGAT1 or DGAT2 α (as assessed by immunoblotting with an anti-FLAG antibody) were assayed at low magnesium concentration using increased amounts of cold oleoyl CoA in the presence of exogenous diacylglycerol. The results are provided in Fig. 3C. Lipids were extracted and separated by TLC and TG accumulation was visualized by charring and quantified by densitometry.

Example 4: Analysis of DGAT2 α mRNA expression in various tissues and in adipocyte differentiation

The tissue distribution of human DGAT2 α mRNA was analyzed. The results are provided in Fig. 4.

DGAT2 α expression increases during 3T3-L1 adipocyte differentiation. Mouse 3T3-L1 adipocyte differentiation was induced and mRNA were isolated at the indicated times shown in Fig. 5. Quantitation of DGAT2 α mRNA levels in triplicate samples was performed by Phosphorimager analysis and corrected for loading relative to actin expression. The results are shown in Fig. 5.

Summary of DGAT2 α

- mouse DGAT2 α has no sequence homology to DGAT1
- mouse DGAT2 α diacylglycerol acyltransferase activity inhibited by high magnesium concentrations;
- human DGAT2 α RNA expression in many tissues, highest levels found in liver, adipose tissue, and mammary gland
- mouse DGAT2 α markedly increased mRNA expression during 3T3-L1 adipocyte differentiation.

Example 5: Characterization of MGAT1 enzymatic activity

The two major pathways for synthesizing diacylglycerol are shown in Figure 9.

Figure 9 depicts the roles of DGAT and monoacylglycerol acyltransferases (MGAT) in the synthesis of triacylglycerides. The examples above describe the activity of DGAT2 α . In the following section, the activity of MGAT1 is described.

MATERIALS AND METHODS

Cloning of MGAT1 cDNA. Mouse expressed sequence tags (ESTs) encoding MGAT1 were identified by BLAST database searches through their sequence homology to DGAT2 from *Mortierella rammaniana* (accession no. AF391089). Based on these ESTs, primers were designed to amplify the complete coding sequence of MGAT1 from mouse liver RNA by reverse transcription (SuperScript Choice System, Gibco BRL, Rockville, Maryland) and PCR (Takara *Ex Taq*, Panvera, Madison, Wisconsin). The MGAT1 sequence has been deposited in GenBank (accession no. AF384162).

Insect Cell Expression Studies. MGAT1 was tagged with an N-terminal FLAG epitope (MGDYKDDDDG, epitope underlined; SEQ ID NO:17) and expressed in *Spodoptera frugiperda* Sf9 insect cells as described. MGAT1 without FLAG was also expressed to determine whether the presence of FLAG, which permits the detection and assessment of expression levels, alters MGAT1 activity. Briefly, the MGAT1 coding sequence (with or without FLAG) was subcloned into pVL1393 baculovirus transfer vector (PharMingen, San Diego, CA). Recombinant baculoviruses were generated by cotransfecting Sf9 insect cells with the transfer vector and BaculoGold DNA (PharMingen). High-titer viruses used for MGAT1 expression were obtained after two rounds of amplification. FLAG-tagged-DGAT1 (accession no. AF078752) and -DGAT2 (accession no. AF384160) were also expressed as controls. To prepare membrane fractions, cells were typically infected with virus for 3 days, washed with PBS, and homogenized by 10 passages through a 27-gauge needle in 1 mM EDTA, 200 mM sucrose, 100 mM Tris-HCl, pH 7.4. Total membrane fractions (100,000 x g pellet) were resuspended in homogenization buffer and frozen at -80°C until use. Expression of MGAT1, DGAT2, and FLAG-tagged proteins in 5 μ g of membrane protein was verified by immunoblotting with an antiserum raised against the C-terminus (amino acids 295-316) of MGAT1, an antiserum against DGAT2, and an anti-Flag M2 antibody (Sigma, St. Louis, Missouri), respectively.

In Vitro Acyltransferase Assays. Generally, acyltransferase activities were assayed under apparent V_{\max} conditions for 5 min in a final volume of 200 μ l. Each reaction contained 100 μ g of membrane proteins, 5 mM $MgCl_2$, 1.25 mg/ml bovine serum albumin (BSA), 200 mM sucrose, 100 mM Tris-HCl, pH 7.4, 25 μ M acyl donor, and 200 μ M acyl acceptor. Non-polar acyl acceptors (diacylglycerol, monoacylglycerol, cholesterol, phosphatidic acid, and sphingosine) were dispersed as phosphatidylcholine liposomes (molar ratio \approx 0.2), and polar acyl acceptors (glycerol-3-phosphate, dihydroxyacetone phosphate, lysophosphatidic acid, and lysophosphatidylcholine) were dissolved in water. Reactions were started by adding protein and terminated by adding 4 ml of chloroform:methanol (2:1 v/v). The extracted lipids were dried, separated by TLC with hexane:ethyl ether:acetic acid (80:20:1 v/v/v), visualized with iodine vapor, and identified with lipid standards. For experiments with radiolabeled substrates, TLC plates were exposed to x-ray film to assess the incorporation of radioactivity into lipid products.

Specifically, DGAT activity was measured as described. Cases et al. (1998) *Proc. Natl. Acad. Sci. USA* 95:13018-13023. MGAT activity was determined by measuring the incorporation of the [^{14}C]oleoyl moiety into diacylglycerol with 25 μ M [^{14}C]oleoyl CoA (specific activity, \sim 20,000 dpm/nmol) and 200 μ M exogenously added *sn*-2 monooleoylglycerol. In some assays, [^{14}C]*sn*-1 monooleoylglycerol (specific activity, 18 μ Ci/ μ mol, 200 μ M final concentration, American Radiolabeled Chemicals, St. Louis, Missouri) was used as a radiolabeled tracer to measure MGAT activity in the presence of unlabeled oleoyl CoA (25 μ M). The dependence of MGAT1 activity on monoacylglycerol and fatty acyl CoA as substrates was determined by assaying MGAT1 with various concentrations of oleoyl CoA or monooleoylglycerol in the presence of 400 μ M monooleoylglycerol or 50 μ M oleoyl CoA, respectively. Diacylglycerol mass was quantified by densitometry after the lipid products were separated by TLC and visualized by immersing the plate in a solution of 10% cupric sulfate and 8% phosphoric acid and heating at 180°C for 30 min. Stereoisomers of monoacylglycerol (*sn*-1-, *sn*-2-monooleoylglycerol, and 3-monostearoylglycerol) were from Sigma. MGAT activity in tissues was measured in particulate fractions prepared from pooled tissues of three 15-week-old male mice.

Mammalian Cell Expression Studies. For mammalian cell expression, FLAG-tagged MGAT1 was subcloned into a pcDNA3 vector and transfected into COS-7 or CHO

cells with Fugene 6 (Roche Diagnostics, Chicago, Illinois). FLAG-tagged-DGAT1, -DGAT2, and -ACAT2 (cholesterol acyltransferase, accession no. AF078751) were expressed as controls. Membrane fractions were prepared as described for insect cells. Expression of FLAG-tagged proteins (in 20 µg of membrane proteins) was verified by immunoblotting with the anti-FLAG M2 antibody. For immunocytochemistry, cells were grown and transfected on glass cover slips. Two days after transfection, cells were fixed in acetone:methanol (1:1) for 2 min and incubated in PBS containing 3% BSA and 0.2% Triton X-100 for 1 h at room temperature. Samples were then incubated sequentially with 4 µg/ml anti-FLAG antibody (Sigma) for 1 h and 10 µg/ml FITC-conjugated goat anti-mouse IgG (CalBiochem, Pasadena, California) for 30 min. Antibodies were diluted in PBS containing 3% BSA and 0.02% Triton X-100. MGAT activities in membranes of transfected cells were assayed as described above.

MGAT1 Tissue Expression Pattern in Mice. To determine tissue distribution of MGAT1 expression, a mouse multiple tissue blot (SeeGene, Seoul, Korea), a blot of total RNA from indicated tissues, and a poly-A⁺ RNA blot (CLONTECH, Palo Alto, California) were hybridized with ³²P-labeled probes generated by random priming (Amersham) with MGAT1 cDNA as the template.

Results

Identification of an MGAT1 Gene. Mouse MGAT1 cDNA was originally identified as a DGAT candidate (DC) gene through its homology to genes encoding DGAT2. Mouse MGAT1 is also referred to herein, and in Cases et al. ((2001) *J. Biol. Chem.* 276:38870-38876), as mouse DC2. The amino acid sequence of mouse MGAT1 is set forth in SEQ ID NO:06; and the nucleotide sequence encoding mouse MGAT1 is set forth in SEQ ID NO:05. The open reading frame of the MGAT1 cDNA predicts a 335-amino acid protein, which is 40% identical to mouse DGAT2, as shown in Figure 10A, and has a predicted molecular mass of 38.8 kDa. Like DGAT2, MGAT1 contains sequences similar to a domain of phosphate acyltransferases. MGAT1 also possesses two putative N-linked glycosylation sites and a potential tyrosine phosphorylation site. The hydrophobicity plot for MGAT1 is similar to that for DGAT2 and predicts at least one transmembrane domain (amino acids 21-43) in the amino terminus, as shown in Figure 10B. Sequences for the human MGAT1 homologue have been reported as human DC2, a member of the DGAT2 gene family. The mouse MGAT1 gene is located on chromosome 1 (accession no. AC079223) and its human homologue is on chromosome 2 (accession no. NT_005126).

Mouse MGAT1 expressed in Insect Cells. To examine the biochemical activity of MGAT1 protein, we expressed FLAG epitope-tagged and non-tagged versions of the cDNA in insect cells. The non-FLAG-tagged version migrated on SDS-PAGE with an apparent molecular mass of ~33 kDa. As expected, the FLAG-tagged version migrated slightly more slowly because of the FLAG epitope. Because MGAT1 shares sequence homology with DGAT2, we first examined whether membranes expressing MGAT1 have DGAT activity. With either [¹⁴C]dioleoylglycerol or [¹⁴C]oleoyl CoA as a radiolabeled substrate, MGAT1-expressing membranes incorporated more radioactivity into triacylglycerols than membranes expressing wild-type viral proteins or heat-inactivated MGAT1 indicating that these membranes have DGAT activity. However, this DGAT activity was significantly less than that in control membranes expressing DGAT2, even though MGAT1 protein was expressed at a higher level.

Since the assays with [¹⁴C]oleoyl CoA radiolabel revealed that a significant amount of [¹⁴C]oleoyl CoA was incorporated into diacylglycerol in membranes expressing MGAT1, we suspected that MGAT1 possesses MGAT activity. To test this possibility, membranes expressing MGAT1 were assayed with either [¹⁴C]monooleoylglycerol or [¹⁴C]oleoyl CoA as the labeled substrate. In both cases, membranes expressing MGAT1 catalyzed the incorporation of the label into diacylglycerol, establishing that the MGAT1 protein possesses MGAT activity. This MGAT activity was confirmed by its dependence on MGAT substrates; when unlabeled MGAT substrate (monooleoylglycerol or oleoyl CoA) was provided over a range of concentrations while the other substrate was held constant, the mass of diacylglycerol synthesized was dependent on the concentration of substrate, as shown in Figure 11. Further, the acyltransferase activity of MGAT1 appeared to be specific for monoacylglycerol (and possibly diacylglycerol) as the acyl group acceptor; no acyltransferase activity was found in MGAT1-expressing membranes when glycerol-3-phosphate, dihydroxyacetone phosphate, lysophosphatidate, lysophosphatidylcholine, sphingosine, or cholesterol were used as the [¹⁴C]oleoyl CoA acceptor.

Next, since there are three stereoisomers of monoacylglycerol, we determined whether MGAT1 can acylate each of these stereoisomers. In these assays [¹⁴C]oleoyl CoA was used as the acyl donor and either *sn*-1-monooleoylglycerol, *sn*-2-monooleoylglycerol, or *sn*-3-monostearoylglycerol as the acyl acceptor. When *sn*-1-monooleoylglycerol or *sn*-3-monostearoylglycerol was used, the major product was *sn*-1,3- diacylglycerol. On the other

hand, when *sn*-2-monooleoylglycerol was used, the major product was *sn*-1,2(2,3)-diacylglycerol. These results indicate that MGAT1 can acylate each of the stereoisomers of monoacylglycerol, mainly at the *sn*-1 or *sn*-3 position. We also found that the specific activities of MGAT1 using either *sn*-1-monooleoylglycerol or *sn*-2-monooleoylglycerol as substrates were similar and that both increased proportionally with MGAT1 protein levels, as shown in Figure 12. These findings indicate that MGAT1 expressed in insect cells can use *sn*-1-monooleoylglycerol and *sn*-2-monooleoylglycerol equally well as substrates *in vitro*.

Mouse MGAT1 expressed in Mammalian Cells. To examine whether MGAT1 protein expressed in mammalian cells also has MGAT activity, we expressed mouse MGAT1 cDNA and control cDNAs transiently in monkey kidney COS-7 cells, as shown in Figure 13. In cells expressing MGAT1, immunofluorescence microscopy demonstrated a perinuclear and reticular staining pattern, consistent with distribution of the protein in the endoplasmic reticulum. COS-7 cell membranes expressing MGAT1 incorporated more radioactivity into diacylglycerol when [¹⁴C]oleoyl CoA and *sn*-2-monooleoylglycerol were provided as substrates, indicating that these membranes possess MGAT activity. Similar levels of MGAT activity were found when *sn*-1-monooleoylglycerol was used as the acyl acceptor. In contrast, MGAT activity was not detected in control membranes, except in those expressing DGAT1, which appeared to possess a low level of MGAT activity. Similar results were found when these cDNAs were expressed in Chinese hamster ovary (CHO) cells.

Tissue Distribution of Mouse MGAT1 Expression. MGAT1 mRNA expression was highest in the stomach and kidney; lower levels of expression were present in white and brown adipose tissue, uterus, and liver. MGAT1 mRNA expression was not detected in the small intestine. Because MGAT activity had not previously been reported in mice, we performed MGAT assays on membranes from mouse tissues. As demonstrated for many other species, the highest activity was found in the small intestine, as shown in Figure 14. MGAT activity was also detected at significant levels in the stomach, kidney, adipose tissue, and liver, where MGAT1 is expressed.

It is apparent from the above results and discussion that polynucleotides encoding mammalian DGAT2 α and MGAT1 enzymes, as well as novel polypeptides encoded thereby, are provided. The subject invention is important for both research and therapeutic applications. Using the DGAT2 α probes of the subject invention, the role of DGAT2 α and

its regulation in a number of physiological processes can be studied *in vivo*. The subject invention also provides for important new ways of treating diseases associated with DGAT2 α and MGAT, such as hypertriglycemia and obesity, as well as in the production of tryglycerides.

5

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. The citation of any publication is for its disclosure prior to the filing date and should not be construed as an admission that the present invention is not entitled to antedate such publication by virtue of prior invention.

10

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it is readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

15

CLAIMS

What is claimed is:

1. A mammalian polynucleotide present in other than its natural environment
5 encoding a polypeptide that exhibits monoacylglycerol and/or diacylglycerol transferase activity and comprising a nucleotide sequence that has at least 50% nucleotide sequence identity to a sequence selected from the group consisting of SEQ ID NO:01, 03, 05, 07, 09, 11, 13, 15, and 18.

10 2. The polynucleotide according to claim 1, wherein said encoded polypeptide is DGAT2 α .

3. The polynucleotide according to claim 1, wherein said encoded polypeptide is MGAT1.
15

4. A mammalian polypeptide present in other than its naturally occurring environment, wherein said polypeptide is selected from the group consisting of diacylglycerol acyltransferase 2 α (DGAT2 α) and monoacylglycerol acyltransferase-1 (MGAT1).
20

5. The polypeptide according to Claim 4, wherein said polypeptide has an amino acid sequence that is substantially the same as or identical to a sequence selected from the group consisting of SEQ ID NO:02, SEQ ID NO:04, SEQ ID NO:06, SEQ ID NO:08, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, and SEQ ID NO:22.
25

6. The polypeptide according to claim 4, wherein said polypeptide is substantially pure.

30 7. An expression cassette comprising a transcriptional initiation region functional in an expression host, a polynucleotide having a nucleotide sequence found in the nucleic acid according to claim 1 under the transcriptional regulation of said transcriptional initiation region, and a transcriptional termination region functional in said expression host.

8. A cell comprising an expression cassette according to claim 7 as part of an extrachromosomal element or integrated into the genome of a host cell as a result of introduction of said expression cassette into said host cell.

5

9. The cellular progeny of the cell according to claim 8.

10. A method of producing a DGAT2 α or an MGAT1 polypeptide, said method comprising:

10 growing a cell according to claim 8, whereby said polypeptide is expressed; and
isolating said polypeptide substantially free of other proteins.

11. A monoclonal antibody binding specifically to a DGAT2 α or an MGAT1 polypeptide.

15

12. The monoclonal antibody according to claim 11, wherein said antibody binds specifically to DGAT2 α and inhibits diglyceride acyltransferase activity of said polypeptide.

13. The monoclonal antibody according to claim 11, wherein said antibody binds specifically to MGAT1 and inhibits monoacylglycerol acyltransferase activity of said polypeptide.

20

14. The monoclonal antibody according to Claim 11, wherein said antibody is a humanized antibody.

25

15. A method for inhibiting the activity of a protein according to claim 4, said method comprising:
contacting said protein with an agent that inhibits the acyltransferase activity of said protein.

30

16. The method according to Claim 15, wherein said agent is a small molecule.

17. The method according to Claim 15, wherein said agent is an antibody.

18. The method according to Claim 17, wherein said agent is a monoclonal antibody.

5 19. A method of modulating a symptom in a mammalian host of a disease condition associated with the acyltransferase activity of a DGAT2 α or an MGAT1 protein, said method comprising:

administering to said host a pharmaceutical composition comprising an effective amount of an active agent that modulates said DGAT2 α or MGAT1 activity in said host.

10 20. The method according to claim 19, wherein said symptom is hypertriglycemia.

21. The method according to claim 19, wherein said symptom is obesity.

15 22. A method of producing a triacylglycerol, said method comprising:
contacting a diacylglycerol and fatty acyl CoA with a DGAT2 α polypeptide under conditions sufficient to said triacylglycerol to be produced.


20 23. A method of identifying an agent that inhibits an acyltransferase activity of a DGAT2 α or an MGAT1 polypeptide, the method comprising:

contacting said DGAT2 α or MGAT1 polypeptide with a test agent in the presence of magnesium ions, a fatty acyl CoA, and an acyl acceptor; and

25 determining the effect, if any, of the test agent on the production of acylated acceptor.


1 / 20


FIG. 1



2 / 20

FIG. 2





4 / 20

FIG. 3B
B. DGAT2 α activity as a function of
Diacylglycerol






FIG. 3C
C. DGAT2 α activity as a function of Oleoyl
Coenzyme A




5 / 20

FIG. 4

A. Tissue distribution of human DGAT2 α 

6 / 20

FIG. 5



MKTLIAAYSGVLGRERAELPAAKNKNKGSALSREGSGRWGTGSSILSALQDIFSVTWLNRSKVEKQLQ
 VISVLQWVLSFLVLGVACSVILMYTFCTDCLWLAIVLYPTWLAFDWNTPKKGRRSQWVRNWAIVRYFRD
 YFPIQLVKTNLLTTRNYIFGYHPHIGLGAFCNFSTEATEVSKFPGRPYLATLAGFRMPVLREY
 LMSGGICLVNRDITDYLKNGSGNAIIIVGGAAEELSMPGNVATLNRKGFVKLALRHGADLVPT
 YSFGENEVYKQVIFEEGSGRWVKFKQYIGFAPCIFHGRGLFSDDTWGLVPYSKPITTVVGEPIITVPK
 LEHPTQKQIDILYHAMVMEALVKLFDNHKTGFLPETEVLEWN (SEQ ID NO: 04)

FIG. 6B

ATGAAGACCCTCATCGCGCCTACTCCGGGGTGAGCTGGGGCGGAAGCTGCCCCGAG
GAAAAACAAGAAATAAGGATCTGCCCTGTACGCGAGGGGTCTGGGCGATGGGGCACTGGGCTCCAGCATC
CTCTCAGCCCTCCAAGACATCTTCTCTGTCACTGGCTCAACAGATCTAAGGTGGA AAAACAGCTGCAG
GTCACTCTCAGTACTACAATGGGTCTTATCTTCTGGTGTAGGAGTGGCTGCAGTGTCTATCTCTCATG
TACACCTTCTGCAACAGACTGTGGGTGATAGCTGTGTCTACTTCACTGGCTGGCACTTTGACTTGAAC
AGCCCCAAGAAAGTGGCAGAGATCGCAGTGGGTGGAAACTGGGCCGTGTGGCGTACTTCCGAGAC
TACTTTCCATCCAGCTGGTGAAGACACAAACCTGTGACCACCAAGAACTATATCTTTGGGATACCA
CCCCATGGCATACTGGGCCCTGGGTGGCTTCTGTAACTTCAGCACAGAGCTACTGAAGTCAGCAAGAAG
TTTCTCGGCATAAGGCCCTATTGGCTACGTTGGCYGGTAACTTCCGGATGCCCTGTGCTTCGCGAGTAC
CTEATGTCTGGAGGCATCTGCCCTGTCAACCGAGACACCATAGACTACTTGTCTCTCCAAGAAATGGGAGT
GGCAATGCTATCATCATCGTGGTGGGAGTGCAGCTGAGTCCCTGAGCTCCATGCTGGCAAGAACCGCA
GTCACCCCTGAAGAACCGCAAGGCTTTGTGAAGCTGGCCCCGTGCCCATGGAGCTGATCTGGTTCTCCACT
TATTCTTTGGAGAGAATGAGGTATACAAGCAGGTGATCTTTGAGGAGGGTTCTTGGGGCCGATGGGTC
CAGAAGAAGTTCCAGAAGTATATTGGTTTCGCCCTGTCATCTTCCATGGCCGAGGCCCTCTTCTCTCT
GACACTGGGGGTGGTGGCTTACTTCAAGCCCCATCACACCGTCTGGGGGAGGCCCATCACTGTCTCCC
AAGCTGGAGCACCCGACCCAGAAAGACATCGACTGTACCATGCTGTACATGGAGGCCCTGGTGAAG
CTCTTTGACAATCACAAAGACCAAAATTTGGCCCTNCCAGAGACTGAGGTGCTGGAGGTGAACCTGA

(SEQ ID NO: 03)

8 / 20

FIG. 7A**Human DAGT2 α amino acid sequence**

MKTLIAAYSGVLRGERQAEADRSQRSHGGPALSREGSGRWGTGSSILSALQDLFSVTWLNRSKVEKQLQV
ISVLQWVLSFLVLGVACSAILMYIFCTDCWLI AVL YFTWL VFDWNTPKKGRRSQWVRNWAVWRYFRDYF
PIQLVKTHNLLTTRNYIFGYHPHGIMGLGAF CNFSTEATEVSKKFPGIRPYLATLAGNFRMPVLREYLS
GGICPVSRDTIDYLLSKNGSGNAIIIVVGGAAESLSSMPGKNAVTLRNRKGFVKLALRHGADLVPIYSFG
ENEVYKQVIFBEGSWGRWVQKKFKYIGFAPCIFHGRGLFSSDTWGLVPYSKPITTVVGEPITIPKLEHP
TQQDIDLYHTMYMEALVKLFDKHKTKFGLPETEVLEV N (SEQ ID NO:02)

FIG. 7B**Human DAGT2 α nucleic acid sequence**

TTACGCCATGAAGACCCTCATAGCCGCCTACTCCGGGGTCTGCGCGCGGAGCGTCAGGCCGAGGCTGACCG
GAGCCAGCGCTCTCACGGAGGACCCGTGTGCGCGGAGGGGTCTGGGAGATGGGGCACTGGATCCAGCATCCT
CTCCGCCCTCCAGGACCTCTTCTGTGTCACCTGGCTCAATAGGTCCAAGGTGGAAAAGCAGCTACAGGTCAT
CTCAGTGCTCCAGTGGGTCTGTCTCTCTTGTACTGGGAGTGGCCTGCAGTGCCATCCTCATGTACATATT
CTGCACTGATTGCTGGCTCATCGCTGTGCTCTACTTCACTTGGCTGGTGTTTGACTGGAACACACCCAAGAA
AGGTGGCAGGAGGTACAGTGGGTCCGAAACTGGGCTGTGTGGCGCTACTTTGAGACTACTTTCCCATCCA
GCTGGTGAAGACACACAACCTGCTGACCACAGGAAGTATATCTTTGGATACCAACCCCATGGTATCATGGG
CCTGGGTGCCTTTCTGCAACTTCAGCACAGAGGCCACAGAAGTGAGCAAGAAGTCCCAGGCATACGGCCTTA
CCTGGCTACACTGGCAGGCAACTTCCGAATGCCTGTGTTGAGGGAGTACCTGATGTCTGGAGGTATCTGCCC
TGTCAGCCGGGACACCATAGACTATTTGCTTTCAAAGAATGGGAGTGGCAATGCTATCATCATCGTGGTCGG
GGGTGCGGCTGAGTCTCTGAGCTCCATGCCTGGCAAGAATGCAGTCACCCCTGCGGAACCGCAAGGGCTTTGT
GAAACTGGCCCTGCGTCATGGAGCTGACCTGGTTCCTATCTACTCCTTTGGAGAGAATGAAGTGTACAAGCA
GGTGATCTTCGAGGAGGGCTCCTGGGGCGATGGGTCCAGAAGAAGTTCAGAAATACATTGGTTTTCGCCCC
ATGCATCTTCCATGGTCGAGGCCTCTTCTCCTCCGACACCTGGGGGCTGGTGCCCTACTCCAAGCCCATCAC
CACTGTTGTGGGAGAGCCCATCACCATCCCCAAGCTGGAGCACCCAACCCAGCAAGACATCGACCTGTACCA
CACCATGTACATGGAGGCCCTGGTGAAGCTCTTCGACAAGCACAAAGACCAAGTTCGGCCTCCCGGAGACTGA
GGTCCTGGAGGTGAACTGAGCCAGCCTTCGGGGCCAATTCCCTGGAGGAACCAAGCTGCAAATCACTTTTTTG
CTCTGTA (SEQ ID NO:01)

9 / 20

FIG. 8A**Mouse DC2 amino acid sequence**

MMVEFAPLNTPLARCLQTAAVLQWVLSFLLLVQVCIGIMVMLVLYNYWFLYIPYLWVFYYDWRTPEQGGRR
RWNWVQSWPVWKYFKEYFPICLVKTQDLDPGHNYIFGFHPHGIFVPGAFAFNFTKYSDFKKLFPGFSTSYL
HVAKIWFCFPLFREYLSMNGPVSVSKESSLHVLSKDGGSNVSIIVLGGAKEALEAHPGTFTLCIRQRKGF
VKMALTHGASLVPVFSFGENDLYKQINNPKGSWLRITQDAMYDSMGVALPLIYARGIFQHYFGIMPYRKL
IYTVVGRPIPVQQILNPTSEQIEELHQTYLEELKKLFENEHKGKYGIPHEHETLVFK (SEQ ID NO:06)

Mouse DC2 nucleic acid sequence

ATGATGGTCGAGTTCGCGCCACTCAACACCCGCTGGCACGGTGCCTACAGACCGCTGCGGTGCTGCAGTGGGT
CCTGTCCTTCCTCCTGCTCGTGCAGGTGTGCATTGGAATTATGGTGATGCTGGTCTGTACAACTATTGGTTCC
TTTACATCCCATATCTGGTCTGGTTTTACTATGACTGGAGAACCCAGAGCAAGGAGGCAGAAGATGGAAGTGG
GTCCAAAGCTGGCCTGTGTGGAAGTATTTAAGGAGTATTTTCCAATCTGTCTTGTCAAACCGCAGGATTTGGA
TCCGGGTACAAATTATATATTTGGGTTTCACCCCTCATGGAATATTCGTGCCTGGAGCCTTTGGAAATTTTGT
CAAAATACTCGGACTTCAAGAAGCTATTTCTGGCTTTACATCGTATCTCCACGTGGCCAAGATCTGGTTCTGT
TTCCCGTTGTTCCGAGAATATCTGATGAGTAACGGGCGGTTTCAGTCTCTAAGGAGAGTTTGTCTCATGTGCT
GAGCAAGGATGGAGGTGGCAATGTCTCAATCATTTGCTCCTCGGAGGTGCAAGGAGGCGCTGGAGGCTCACCCAG
GAACATTCACCTGTGCATCCGCCAGCGCAAAGGGTTTGTAAAGATGGCCTTGACCCATGGTGCCAGTTTGGTT
CCAGTATTTTCTTTTGGTGAAAATGATCTATATAAGCAAATTAACAACCCCAAAGGCTCCTGGCTACGAAGTAT
ACAAGACGCAATGATGATTCAATGGGAGTAGCCTTGCCACTGATATATGCCAGAGGAATTTCCAGCACTACT
TTGGCATAATGCCCTATCGGAAGCTGATCTACACTGTTGTTGGCCGCCCTATCCCTGTTTCAAGCAGATTCTGAAC
CCGACCTCAGAGCAGATTGAAGAGCTGCATCAGACATACCTAGAGGAGCTAAAGAACTATTCAATGAACACAA
AGGGAAATATGGGATTCCGGAGCACGAACTCTGGTATTTAAATAA (SEQ ID NO:05)

Human DC2 amino acid sequence

MKVEFAPLINIQLARRLQTVAVLQWVLSFLTGPMSIGITVMLIIHNYLFLYIPYLMWLYFDWHTPERGGRR
SSWIKNWTLLWKHFKDYFPIHLIKTQDLDP SHNYIFGFHPHGIMAVGAFGNFSVNYSDFKDLFPGFSTSYLH
VLPLWFWCPVFRFYVMSVGLVSVSKSVSYMVSKEGGNISVIVLGGAKESLDAHPGKFTLFIQRQRKGFV
KIALTHGASLVPVVSFGENELFKQTDNPEGSWIRTVQNKLQKIMGFALPLFHARGVFQYNFGIMTYRKAI
HTVVGRPIPVROTINPTQEQIEELHQTYMEELRKLFEHKGKYGIPHEHETLVLK (SEQ ID NO:08)

Human DC2 nucleic acid sequence

CGTGGGTGCAGGCTGCAGTGGCTGGCGCCGCTCCTCGCCCGGCCAGGCCATGAAGGTAGAGTTTGCACCGCTCAA
CATCCAGCTGGCGCGGCGGCTGCAGACGGTGGCCGTGCTGCAGTGGGTCTTTTCTTTTCTTACAGGGCCGATGT
CCATTGGAATCACTGTGATGCTGATCATAACAATACTATTTGTTTCTTTTACATCCCTTATTTGATGTGGCTTTAC
TTTGACTGGCATAACCCAGAGCGAGGAGGCAGGAGATCCAGCTGGATCAAAAATTGGACTCTTTGGAAACACTT
TAAGGACTATTTTCCAATTCATCTTATCAAACTCAAGATTTGGATCCAAGTCACAATAATATATTTGGGTTTC
ACCCCATGGAATAATGCGAGTTGGAGCCTTTGGGAATTTTCTGTAAATTATCTGACTTCAAGGACCTGTTT
CCTGGCTTTACTTTCATATCTTCACGTGCTGCCACTTTGGTTCTGGTGTCTGTCTTTGAGAATATGTGATGAG
TGTGGGCTGGTTTTCAGTTTCCAAGAAAAGTGTGTCTTACATGGTAAGCAAGGAGGGAGGTGGAAACATCTCTG
TCATTGTCTTGGGGGTGCAAAAGAATCACTGGATGCTCATCCTGGAAAGTTCACTCTGTTTCATCCGCCAGCGG
AAAGGATTTGTTAAATTTGCTTTGACCCATGGCGCCTCTCTGGTCCCAGTGGTTTCTTTTGGTGAAAATGAAC
GTTTAAACAACTGACAACCTGAAGGATCATGGATTAGAAGTGTTCAGAATAAACTGCAGAAGATCATGGGGT
TTGCTTTGCCCTGTTTCATGCCAGGGGAGTTTTTTCAGTACAATTTTGGCCTAATGACCTATAGGAAAGCCATC
CACACTGTTGTTGGCCGCCGATCCCTGTTTCGTGAGACTCTGAACCCGACCCAGGAGCAGATTGAGGAGTTACA
TCAGACCTATATGGAGGAAGTTAGGAAATGTTTGGAGAACACAAAGGAAAGTATGGCATTCCAGAGCAGAGA
CTCTGTTTTAAATGACTTGACTATAAAAAAATAAAAAATAAAATGACTTGGCTGTAATAAGGCA
TAAAGAAGGATAAGAGACC (SEQ ID NO:07)

10 / 20

FIG. 8B

Mouse DC3 amino acid sequence (partial sequence, about 100 amino acids gap in the center of the protein)

MKTEHLQSLSLQWPLSYVAMFWIVQPLLCILLETPLWPLPTVYFVWLLLDWKTPDKGRRSDWVRNWNV
WNHIRDYFPITILKTKDLSPSENYIMGVHPHGLLTFGAFCNFCTEATGFSKTFPGITPHLATLSWFFKIP
IIRDYIMAKGLCSVSQASIDYLLSHGTGNLVGIPITTVGEALPLPQVKNPSPHIVDKYHALYMDALYKL
FEQHKVQYGCSTNTQKLIFL (SEQ ID NO:10)

Mouse DC3 nucleic acid sequence (partial sequence, about 100 amino acids gap in the center of the protein)

TTACCTGCCTCAGGGTCCTGGGCATCATGTCTTGCTCTATGAAGACTGAACACTTACAGAGTCTGAGCCTTCTG
CAGTGGCCCTTGAGCTACGTTGCCATGTTTTGGATTGTGCAGCCATTGTTAATTTGCCTATTGTTACACCCTT
GTGGCCGCTACCAACAGTTTACTTTGCTGTTACTTCTCGACTGGAAGACTCCAGATAAAGGTGGCAGGCGTT
CAGACTGGGTACGGAAGTGAATGTCTGGAACCATCAGGGACTATTTCCCCATTACAATCCTGAAGACTAAG
GACCTGTCACCTTCAAGAACTACATCATGGGGTCCACCCCATNGGTCTCCTGACCTTCGGTGCCTTCTGCAA
CTTCTGCACTGAGGCCACAGGCTTCTCGAAGACCTTCCAGGCATCACTCCTCACTTGGCCACAC (SEQ ID
NO:09)

Human DC3 amino acid sequence

MAFFSRLNLQEGLOTFFVLQWIPVYIFLVWILQPLFVYLLETSLWPLPVLYFAWLFLDWKTPERGGRRSA
WVRNWCWVTHIRDYFPITILKTKDLSPHENYLMGVHPHGLLTFGAFCNFCTEATGFSKTFPGITPHLATL
SWFFKIPFVREYLMAGVCSVSQPAINYLLSHGTGNLVGIVVGGVGEALQSVNNTTLLQKRKGFVRTA
LQHGAYLVPSYSFGENEVFNQETFPGETWLRFLQRTFQDTFKILGLNFCTFHGRGFTRGSWGFLPFNRP
ITTVVGEPLPIPRIKRPNQKTVDKYHALYISALRKLFDQHKVEYGLPETQELTIT (SEQ ID NO:12)

Human DC3 nucleic acid sequence

ATCAACTCAGCTTAAGAAGTTTTGGCCTTCTGGTTAGGCTTCTTGCCACAACAGAACAGCACCATAACCATGGC
TTTCTTCTCCCGACTGAATCTCCAGGAGGGCCTCCAAACCTTCTTTGTTTTGCAATGGATCCCAGTCTATATAT
TTTTAGTTTGGATCTTGCAGCCATTGTTCTGTCTACCTGCTGTTTACATCCTTGTTGGCCGCTACCAGTGCTTTAC
TTTGCCTGGTTGTTCTGGACTGGAAGACCCCAAGAGCGAGGTGGCAGGCGTTCCGGCCTGGGTAAAGGAAGTGGT
TGTCTGGACCCACATCAGGGACTATTTCCCCATTACGATCCTGAAGACAAAGGACCTATCACCTGAGCACAAC
ACCTCATGGGGGTTACCCCCATGGCCTCCTGACCTTTGGCGCCTTCTGCAACTTCTGCACTGAGGCCACAGGC
TTCTCGAAGACCTTCCAGGCATCACTCCTCACTTGCCACGCTGTCTGTTCTTCAAGATCCCCTTTGTTAG
GGAGTACCTCATGGCCAAAGGTGTGTGCTCTGTGAGCCAGCCAGCCATCAACTATCTGCTGAGCCATGGCACTG
GCAACCTCGTGGGCATTGTAGTGGGAGGTGTGGGTGAGGCCCTGCAAAGTGTGCCCAACACCACCACCTCATC
CTCCAGAAGCGCAAGGGGTTCTGTGCGCACAGCCCTCCAGCATGGGGCATACCTTGTCCCTTCATATTCCTTTGG
TGAGAACGAAGTTTTCAATCAGGAGACCTTCCCTGAGGGCACGTGGTTAAGGTTGTTCCAAAAAACCTTCCAGG
ACACATTCAAAAAATCCTGGGACTAAATTTCTGTACCTTCCATGGCCGGGGCTTCACTCGCGGATCCTGGGGC
TTCTGCTTTTCAATCGGCCCATTAACACTGTTGTTGGGGAACCCCTTCCAATTCACAGGATTAAGAGGCCAAA
CCAGAAGACAGTAGACAAGTATCACGCACTCTACATCAGTGCCCTGCGCAAGCTCTTTGACCAACACAAAGTTG
AATATGGCCTCCCTGAGACCCAAGAGCTGACAATTACATAACAGGAGCCACATTCCTTATGATCAACCCCCAA
AGCCATGAGGGATCCAAGTAGAGCCACAGAAAAAGAAATTCAGGAGAGGGAAAGATCGTAAGGATGAGAGA
GGAGACCATCCAAGCCAGAAATTATTTAATAAATCAGAGTTCTAGCAATAGAGTCC (SEQ ID NO:11)

11 / 20

FIG. 8C**Human DC4 amino acid sequence**

MLLPSKKDLKTALDVFAVFQWSFSALLITTTVIAVNLYLVVFTPYWVPTVLILTLWLAFDWKTPQRGRRF
TCVRHWRLWKHYSYDFPLKLLKTHDICPSRNYILVCHPHGLFAHGWFGHFATEASGFSKIFPGITPYILT
LGAFWMPFLREYVMSTGACSVSRSSIDFLTHKGTGNMVIVVIGGLAECRYSLPGSSITLVLKNRSGFVR
MALQHGVPPLIPAYAFGETDLYDQHIFTGGFVNRQKWFQSMVHIYPCAFYGRGFTKNSWGLLPYSRPVT
TIVGEPLPMPKIENPSQEIVAKYHTLYIDALRKLFDQHKTKFGISETQELEII (SEQ ID NO:014)

Human DC4 nucleic acid sequence

AATTCGGCTTACTCACTATAGGGCTCGAGCGGGCCCCCGGGCAGGTGCCGACTTCATTTCCAAGTCTGCAC
ACAAATGCAGGCAGTAGCCATGCCTGACAGCCACATGACAGATACTACACCGCTGAATGTGCTCTAACCCCT
GGACTTGGCATTGCCCTACTGTTGAGGAAGCAGTGCGTTTTTCTCCAGTCTTTCAGGTCCCTTCACCAG
GGAACCATTAACCTTGTGCATCAGAACAAGGACATTTCTTACATTCTGCAAACACAGTCCTTTCAGTTT
ACTCTTTTTTTGAGGGGGGGGCGCGGGGAACGGAGTCTCGCTCTGTGCGCCAGGCTGGAGTGCAATGGTG
CAATCTCAGCTCACTGCAACCTCTGCCTCCCAGGTCCAAGCGATTCTCCTGCCTCAGCCTCCCGGGTAGC
CGGGACTACAGGCGCCTGCCACCACGCCCCGGCTAATTTTTGTATTTTAGTAGAGACGAGGTTTCGCCGT
GTTGGCAGGCTGGTCTTGGAACTCCTGACCTCAGGTGATTTACTCGCCTCGGCCTCCCAAAGTGCTGGGA
TTACAGGCATGAGCCACTGTGCCAGTCACAAGTTTTATTTTAGCCATTTTGATAAGTGTGAAGTTCCC
TGATGGCTAATGATGTTCCCTTTTCCATGTGCTCATTTGTCTATGCCAGAGAAGATTTGGAGAGGAG
GACGTGAATTGGAGGAAACTGTTCCAGGATTCACCACTCTGGTGGCCACCGCTGGCTCACTGCCATT
GACCACACTGCAGGCAGAGCCTAGTGCAGTGCTGGAGCAGGGCCAGAGAGGAGAGGGCTTACAGTGTGA
ATTCAGCTCAGCTGGGGAAGAAGACACCTTCCCTTCTAGACCTGAATCGGGTTCCCAAGCAACCACTGTG
ATTGCTGTCAACCTCTACCTGGTGGTGTTCACACCATACTGGCCTGTCACTGTGCTTATTCTTACCTGGC
TGGCTTTTGACTGGAAGACCCCTCAGCGAGGCGGGCCCGGTTTACCTGTGTGAGGCACTGGCGCCTGTG
GAAACACTACAGCGATTATTTCCCTCTCAAGCTTCTGAAGACTCATGACATCTGCCCCAGCCGCAACTAC
ATCCTCGTCTGCCACCTCATGGGCTCTTTGCCCATGGATGGTTTGGCCACTTTGCCACAGAGGCCTCAG
GCTTCTCCAAGATATTTCCCTGGCATCACCCCTTACATACTCACACTGGGAGCCTTTTTCTGGATGCCTTT
CCTCAGAGAATATGTAATGTCTACAGGGGCCTGCTCTGTGAGTCGATCCTCCATTGACTTTCTGCTGACT
CATAAAGGCACAGGCAACATGCTCATTGTGGTGATTGGTGGACTGGCTGAGTGCAGATACAGCCTGCCAG
GTTCTTCTACCCCTGGTGTGAAGAACCGGTCTGGCTTTGTGCGCATGGCCCTTCAGCATGGGGTGCCTCT
AATACCTGCCTATGCCTTTGGGGAGACGGACCTCTATGATCAGCACATTTTCACTCCTGGTGGCTTTGTC
AACCCTTCCAGAACTGGTTCCAGAGCATGGTACACATCTACCTTGTGCTTTCTATGGACGTGGCTTCA
CCAAGAACTCCTGGGGCCTTCTGCCCTATAGTCGGCCTGTAACCACCATCGTCGGGGAGCCTCTACCAAT
GCCCAAGATTGAGAATCCAAGCCAGGAGATCGTGGCTAAATATCACACACTCTATATTGATGCCCTACGT
AAACTGTTTGACCAGCATAAAGACCAAGTTTGGTATCTCAGAGACCCAGGAGCTGGAGATAATTTGACAGA
CATCCCCAGTAGCCTTCACCCTGGCTGGAAGGTATGGATGGACCCAGTGAGA (SEQ ID NO:013)

12 / 20

FIG. 8D**Human DC5 amino acid sequence**

MVEFAPLFVPWERRRLQTLAVLQFVFSFLALGKICTVGFIALLFTRFWLLTVLYAAWWYLDLDRDKPRQGGRHI
QAIRCWTIWKYMKDYFPIQLVKTAELDPSRNYIAGFHPHGVLA VGAFANLCTESTGFSSIFPGIRPHLMML
TLWFRAPFFRDYIMSAGLVTSEKESAAHILNRKGGGNLLGIIVGGAQEALDARPGSF TLLLRNRKGFVRLA
LTHGAPLVPIPSFGENDLFDQIPNSSGSWLRYIQNRLQKIMGISLPLFHGRGVFQYSFGLIPYRRPITTVG
KPIEVQKTLHPSEEEVNQLHQHYIKELCNLFEAHKLKFNIPADQHLEFC (SEQ ID NO:16)

Human DC5 nucleic acid sequence

CCACAGCAGAGCTCACAGAACCTGCGGGAGCCAGGCTGACCCGCCAGCATGGTAGAGTTCGCGCCCTTGTT
TGTGCEGTGGGAGCGCAGGCTGCAGACACTTGCTGTCTTACAGTTTGTCTTCTCCTTCTTGGCACTGGGTA
AGATCTGCACTGTGGGCTTCATAGCCCTCCTGTTTACAAGATTCTGGCTCCTCACTGTCTCTGTATGCGGCC
TGGTGGTATCTGGACCGAGACAAGCCACGGCAGGGGGGCCGGCACATCCAGGCCATCAGGTGCTGGACTAT
ATGGAAGTACATGAAGGACTATTTCCCATCCAGCTGGTCAAGACTGCTGAGCTGGACCCCTCTCGGAAC
ACATTGCGGGCTTCCACCCCATGGAGTCCTGGCAGTCGGAGCCTTTGCCAACCTGTGCACTGAGAGCACA
GGCTTCTCTTCGATCTTCCCCGGTATCCGCCCCATCTGATGATGCTGACCTTGTGGTTCCGGGCCCCCTT
CTTCAGAGATTACATCATGTCTGCAGGGTTGGTCACATCAGAAAAGGAGAGTGCTGCTCACATTCTGAACA
GGAAGGGTGGCGGAAACTTGCTGGGCATCATTGTAGGGGGTGCCAGGAGGCCCTGGATGCCAGGCCTGGA
TCCTTCACGCTGTTACTGCGGAACCGAAAGGGCTTCGTACAGGCTCGCCCTGACACACGGGGCACCCCTGGT
GCCAATCTTCTCCTTCGGGGAGAATGACCTATTTGACCAGATTCCCAACTCTTCTGGCTCCTGGTTACGCT
ATATCCAGAATCGGTTGCAGAAGATCATGGGCATCTCCCTCCCACTCTTTATGGCCGTGGTGTCTTCCAG
TACAGCTTTGGTTTAAATACCCTACCGCCGGCCCATCACCAGTGTGGGGAAGCCCATCGAGGTACAGAAGAC
GCTGCATCCCTCGGAGGAGGAGGTGAACCAGCTGCACCAGCATTATATCAAAGAGCTGTGCAACCTCTTCG
AGGCCCCACAACTTAAGTTCAACATCCCTGCTGACCAGCACTTGGAGTTCTGCTGA (SEQ ID NO:15)

13 / 20

FIG. 9





FIG. 10A

SDOCID: <WO_____02068595A2_1_>

15 / 20

FIG. 10B



16 / 20




FIG. 11

17 / 20





FIG. 12

18 / 20



Western: Anti-Flag

FIG. 13




FIG. 14

20 / 20

FIG. 15A

Human hDC5-284 (SEQ ID NO:19)

MVEFAPLFMPWERRLQTLAVLQFVFSFLALAEICTVGFIALLFTRFWLLTVLY
AAWYLDNRDKPRQGRHIQAIRCWTIWKYMKDYFPISLVKTAELDPSRNYI
AGFHPHGVLAVGAFANLCTESTGFSSIFPGIRPHLMMLTLWFRAPFFRDYIMS
AGLVTSKESAAHILNRKGGGNLLGIIVGGAQEALDARPGSFTLLLRNRKGFV
RLALTHGYQASGKSTLGSVGNWQGFYFGKMAETNADSILVEIFSPFTIKIIF
WCLMPKYLEKFPQRRLSDLRN

FIG. 15B

Human hDC5-334 (SEQ ID NO:20)

MVEFAPLFVPWERRLQTLAVLQFVFSFLALAEICTVGFIALLFTRFWLLTVLY
AAWYLDNRDKPRQGRHIQAIRCWTIWKYMKDYFPISLVKTAELDPSRNYI
AGFHPHGVLAVGAFANLCTESTGFSSIFPGIRPHLMMLTLWFRAPFFRDYIMS
AGLVTSKESAAHILNRKGGGNLLGIIVGGAQEALDARPGSFTLLLRNRKGFV
RLALTHGAPLVPIFSFGENDLFDQIPNSSGSWLR YIQNRLQKIMGISLPLFHGR
GVFQYSFGLIPYRRPITTVVGKPIEVQKTLHPSEEEVNQLHQR YIKELCNLFEA
HKLLKFNIPADQHLFC

FIG. 15C

Mouse mDC5-296 (SEQ ID NO:21)

MVEFAPLLVPWERRLQTFAVLQWVFSFLAALQLCIVIFVGLLFTFRFWLFSVLY
ATWYLDWDKPRQGRPIQFFRRLAIWKYMKDYFPVSLVKTAELDPSRNYI
AGFHPHGVLAAAGAFNLCTESTGFTSLFPGIRSYLMMLTVWFRAPFFRDYIMS
GGLVSSSEKVSADHILSRKGGGNLLAIIVGGAQEALDARPGAYRLLLRNRKGFV
RLALMHGAALVPIFSFGENNLFNQVENTPGTWLRWIQNRLQKIMGISLPLFHG
RGVFQYSFGLMPFRQPITTTIGERRSRAEQDQE

FIG. 15D

SEQUENCE LISTING

<110> Cases, Sylvaine
 Stone, Scot
 Zhou, Ping
 Farese, Robert V.
 Chi-Liang Eric Yen

<120> Mono- and diacylglycerol acyltransferases
 and methods of use thereof

<130> UCAL240WO

<140> Unassigned

<141> 2002-02-21

<150> 60/271,307

<151> 2001-02-23

<150> 09/794,715

<151> 2001-02-26

<150> 10/046,924

<151> 2002-01-14

<160> 22

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1231

<212> DNA

<213> Homo sapiens

<400> 1

ttcagccatg	aagaccctca	tagccgccta	ctccgggggtc	ctgcgcggcg	agcgtcagggc	60
cgaggctgac	cggagccagc	gctctcacgg	aggaccggtg	tcgcgcgagg	ggctctgggag	120
atggggcact	ggatccagca	tcctctccgc	cctccaggac	ctcttctctg	tcacctggct	180
caataggtcc	aaggtggaaa	agcagctaca	ggcatctca	gtgctccagt	gggtcctgtc	240
cttcttcta	ctgggagtg	cctgcagtgc	catcctcatg	tacatattct	gcactgattg	300
ctggctcatc	gctgtgctct	acttcaactg	gctgggtgtt	gactggaaca	cacccaagaa	360
aggtggcagg	aggtcacagt	gggtccgaaa	ctgggctgtg	tggcgctact	ttcgagacta	420
ctttcccatc	cagctgggtga	agacacacaa	cctgctgacc	accaggaact	atatctttgg	480
ataccacccc	catggtatca	tgggcctggg	tgccttctgc	aacttcagca	cagaggccac	540
agaagtgagc	aagaagttcc	caggcatacg	gccttacctg	gctacactgg	caggcaactt	600
ccgaatgcct	gtgttgaggg	agtacctgat	gtctggagg	atctgccttg	tcagccggga	660
caccatagac	tatttgcttt	caaagaatgg	gagtggcaat	gctatcatca	tcgtggctcg	720
gggtgcggct	gagtctctga	gctccatgcc	tggcaagaat	gcagtcaccc	tgcggaaccg	780
caagggcttt	gtgaaactgg	ccctgcgtca	tggagctgac	ctgggttcca	tctactcctt	840
tggagagaat	gaagtgtaca	agcaggtgat	cttcgaggag	ggctcctggg	gccgatgggt	900
ccagaagaag	ttccagaaat	acattgggtt	cgccccatgc	atcttccatg	gtcgaggcct	960
cttctctctc	gacacctggg	ggctgggtgc	ctactccaag	cccatcacca	ctgttggtggg	1020
agagcccatc	accatcccca	agctggagca	cccaaccag	caagacatcg	acctgtacca	1080
caccatgtac	atggaggccc	tgggtgaagct	cttcgacaag	cacaagacca	agttcggcct	1140
cccggagact	gaggtcctgg	aggtgaactg	agccagcctt	cggggccaat	tccttgaggg	1200
aaccagctgc	aaatcacttt	tttgctctgt	a			1231

<210> 2

<211> 388

<212> PRT

<213> Homo sapiens

<400> 2

Met Lys Thr Leu Ile Ala Ala Tyr Ser Gly Val Leu Arg Gly Glu Arg
 1 5 10 15

Gln Ala Glu Ala Asp Arg Ser Gln Arg Ser His Gly Gly Pro Ala Leu
 20 25 30
 Ser Arg Glu Gly Ser Gly Arg Trp Gly Thr Gly Ser Ser Ile Leu Ser
 35 40 45
 Ala Leu Gln Asp Leu Phe Ser Val Thr Trp Leu Asn Arg Ser Lys Val
 50 55 60
 Glu Lys Gln Leu Gln Val Ile Ser Val Leu Gln Trp Val Leu Ser Phe
 65 70 75 80
 Leu Val Leu Gly Val Ala Cys Ser Ala Ile Leu Met Tyr Ile Phe Cys
 85 90 95
 Thr Asp Cys Trp Leu Ile Ala Val Leu Tyr Phe Thr Trp Leu Val Phe
 100 105 110
 Asp Trp Asn Thr Pro Lys Lys Gly Gly Arg Arg Ser Gln Trp Val Arg
 115 120 125
 Asn Trp Ala Val Trp Arg Tyr Phe Arg Asp Tyr Phe Pro Ile Gln Leu
 130 135 140
 Val Lys Thr His Asn Leu Leu Thr Thr Arg Asn Tyr Ile Phe Gly Tyr
 145 150 155 160
 His Pro His Gly Ile Met Gly Leu Gly Ala Phe Cys Asn Phe Ser Thr
 165 170 175
 Glu Ala Thr Glu Val Ser Lys Lys Phe Pro Gly Ile Arg Pro Tyr Leu
 180 185 190
 Ala Thr Leu Ala Gly Asn Phe Arg Met Pro Val Leu Arg Glu Tyr Leu
 195 200 205
 Met Ser Gly Gly Ile Cys Pro Val Ser Arg Asp Thr Ile Asp Tyr Leu
 210 215 220
 Leu Ser Lys Asn Gly Ser Gly Asn Ala Ile Ile Ile Val Val Gly Gly
 225 230 235 240
 Ala Ala Glu Ser Leu Ser Ser Met Pro Gly Lys Asn Ala Val Thr Leu
 245 250 255
 Arg Asn Arg Lys Gly Phe Val Lys Leu Ala Leu Arg His Gly Ala Asp
 260 265 270
 Leu Val Pro Ile Tyr Ser Phe Gly Glu Asn Glu Val Tyr Lys Gln Val
 275 280 285
 Ile Phe Glu Glu Gly Ser Trp Gly Arg Trp Val Gln Lys Lys Phe Gln
 290 295 300
 Lys Tyr Ile Gly Phe Ala Pro Cys Ile Phe His Gly Arg Gly Leu Phe
 305 310 315 320
 Ser Ser Asp Thr Trp Gly Leu Val Pro Tyr Ser Lys Pro Ile Thr Thr
 325 330 335
 Val Val Gly Glu Pro Ile Thr Ile Pro Lys Leu Glu His Pro Thr Gln
 340 345 350
 Gln Asp Ile Asp Leu Tyr His Thr Met Tyr Met Glu Ala Leu Val Lys
 355 360 365
 Leu Phe Asp Lys His Lys Thr Lys Phe Gly Leu Pro Glu Thr Glu Val
 370 375 380
 Leu Glu Val Asn
 385

<210> 3

<211> 1167

<212> DNA

<213> Mus musculus

<220>

<221> misc_feature

<222> (1)...(1167)

<223> n = A,T,C or G

<400> 3

atgaagaccc	tcatacgccgc	ctactdcggg	gtcctgcggg	gtgagcgctcg	ggcggaagct	60
gcccgcagcg	aaaacaagaa	taaaggatct	gccctgtcac	gcgaggggtc	tgggcgatgg	120
ggcaactggct	ccagcatcct	ctcagccctc	caagacatct	tctctgtcac	ctggctcaac	180
agatcyaaagg	tggaaaaaca	gctgcaggtc	atctcagtac	tacaatgggt	cctatccttc	240
ctggtgctag	gagtggcctg	cagtgtcatc	ctcatgtaca	ccttctgcac	agactgctgg	300
ctgatagctg	tgtctactt	cacctggctg	gcatttgact	ggaacacgcc	caagaaaggt	360

```

ggcaggagat cgagtggtt ggcgaactgg gccgtgtggc gctacttccg agactacttt 420
cccatccagc tgggtgaagac acacaacctg ctgaccacca ggaactatat ctttggatac 480
caccceccatg gcatcatggg cctgggtgcc ttctgttaact tcagcacaga ggctactgaa 540
gtcagcaaga agtttctctg cataaggccc tatttggcta cgttggcygg taacttccgg 600
atgcctgtgc ttccgagta cctgatgtct ggaggcatct gccctgtcaa ccgagacacc 660
atagactact tgctctccaa gaatgggagt ggcaatgcta tcatcatcgt ggtgggaggt 720
gcagctgagt ccctgagctc catgcctggc aagaacgcag tcaccctgaa gaaccgcaaa 780
ggctttgtga agctggccct gcgccatgga gctgatctgg ttccactta ttcttttggg 840
gagaatgagg tatacaagca ggtgatcttt gaggagggtt cctggggccg atgggtccag 900
aagaagttcc agaagtatat tggtttccgc cctgcatct tccatggccg aggcctcttc 960
tcctctgaca cctgggggct ggtgcctac tccaagcca tcaccaccgt cgtgggggag 1020
cccatcactg tccccaagct ggagcaccgc acccagaaag acatcgacct gtaccatgcc 1080
atgtacatgg aggccttgg gaagctcttt gacaatcaca agaccaaatt tggcctncca 1140
gagactgagg tgctggaggt gaactga 1167

```

<210> 4

<211> 387

<212> PRT

<213> Mus musculus

<400> 4

```

Met Lys Thr Leu Ile Ala Ala Tyr Ser Gly Val Leu Arg Gly Glu Arg
  1          5          10          15
Arg Ala Glu Leu Pro Ala Ala Lys Asn Lys Asn Lys Gly Ser Ala Leu
          20          25          30
Ser Arg Glu Gly Ser Gly Arg Trp Gly Thr Gly Ser Ser Ile Leu Ser
          35          40          45
Ala Leu Gln Asp Ile Phe Ser Val Thr Trp Leu Asn Arg Ser Lys Val
          50          55          60
Glu Lys Gln Leu Gln Val Ile Ser Val Leu Gln Trp Val Leu Ser Phe
          65          70          75          80
Leu Val Leu Gly Val Ala Cys Ser Val Ile Leu Met Tyr Thr Phe Cys
          85          90          95
Thr Asp Cys Trp Leu Ile Ala Val Leu Tyr Phe Thr Trp Leu Ala Phe
          100          105          110
Asp Trp Asn Thr Pro Lys Lys Gly Gly Arg Arg Ser Gln Trp Val Arg
          115          120          125
Asn Trp Ala Val Trp Arg Tyr Phe Arg Asp Tyr Phe Pro Ile Gln Leu
          130          135          140
Val Lys Thr His Asn Leu Leu Thr Thr Arg Asn Tyr Ile Phe Gly Tyr
          145          150          155          160
His Pro His Gly Ile Met Gly Leu Gly Ala Phe Cys Asn Phe Ser Thr
          165          170          175
Glu Ala Thr Glu Val Ser Lys Lys Phe Pro Gly Ile Arg Pro Tyr Leu
          180          185          190
Ala Thr Leu Ala Gly Asn Phe Arg Met Pro Val Leu Arg Glu Tyr Leu
          195          200          205
Met Ser Gly Gly Ile Cys Leu Val Asn Arg Asp Thr Ile Asp Tyr Leu
          210          215          220
Leu Ser Lys Asn Gly Ser Gly Asn Ala Ile Ile Ile Val Val Gly Gly
          225          230          235          240
Ala Ala Glu Ser Leu Ser Ser Met Pro Gly Lys Asn Ala Val Thr Leu
          245          250          255
Lys Asn Arg Lys Gly Phe Val Lys Leu Ala Leu Arg His Gly Ala Asp
          260          265          270
Leu Val Pro Thr Tyr Ser Phe Gly Glu Asn Glu Val Tyr Lys Gln Val
          275          280          285
Ile Phe Glu Glu Gly Ser Trp Gly Arg Trp Val Lys Lys Phe Gln Lys
          290          295          300
Tyr Ile Gly Phe Ala Pro Cys Ile Phe His Gly Arg Gly Leu Phe Ser
          305          310          315          320
Ser Asp Thr Trp Gly Leu Val Pro Tyr Ser Lys Pro Ile Thr Thr Val
          325          330          335
Val Gly Glu Pro Ile Thr Val Pro Lys Leu Glu His Pro Thr Gln Lys
          340          345          350
Asp Ile Asp Leu Tyr His Ala Met Tyr Met Glu Ala Leu Val Lys Leu

```

355 360 365
 Phe Asp Asn His Lys Thr Lys Phe Gly Leu Pro Glu Thr Glu Val Leu
 370 375 380
 Glu Val Asn
 385

<210> 5
 <211> 1008
 <212> DNA
 <213> Mus musculus

<400> 5
 atgatggctcg agttcgcgcc actcaacacc ccgctggcac ggtgcctaca gaccgctgcg 60
 gtgctgcagt gggtcctgtc cttcctcctg ctcgtgcagg tgtgcattgg aattatgggtg 120
 atgctgggtcc tgtacaacta ttggttcctt tacatcccat atctgggtctg gttttactat 180
 gactggagaa cccagagca aggaggcaga agatggaact ggtccaaag ctggcctgtg 240
 tggaagtatt ttaaggagta ttttccaatc tgtcttgta aaacgcagga ttggatccg 300
 ggtcacaatt atatatattgg gtttcaccct catggaatat tcgtgcctgg agcctttgga 360
 aattttttgta caaaatactc ggacttcaag aagctatttc ctggctttac atcgtatctc 420
 cacgtggcca agatctgggt ctgtttcccg ttgttccgag aatatctgat gagtaacggg 480
 ccggtttcag tgtctaagga gagtttgtct catgtgctga gcaaggatgg aggtggcaat 540
 gtctcaatca ttgtcctcgg aggtgcaaag gaggcgctgg aggctcacc aggaacattc 600
 accctgtgca tccgccagcg caaagggttt gttaagatgg cctgaccca tgggtccagt 660
 ttggttccag tattttcttt tgggtgaaaat gatctatata agcaaattaa caaccccaaa 720
 ggctcctggc tacgaactat acaagacgca atgtatgatt caatgggagt agccttgcca 780
 ctgatatatg ccagaggaat tttccagcac tactttggca taatgcccta tcggaagctg 840
 atctacactg ttgttggccg ccctatccct gttcagcaga ttctgaaccc gacctcagag 900
 cagattgaag agctgcatca gacataccta gaggagctaa agaaactatt caatgaacac 960
 aaagggaaat atgggattcc ggagcacgaa actctggtat ttaataaa 1008

<210> 6
 <211> 335
 <212> PRT
 <213> Mus musculus

<400> 6
 Met Met Val Glu Phe Ala Pro Leu Asn Thr Pro Leu Ala Arg Cys Leu
 1 5 10 15
 Gln Thr Ala Ala Val Leu Gln Trp Val Leu Ser Phe Leu Leu Leu Val
 20 25 30
 Gln Val Cys Ile Gly Ile Met Val Met Leu Val Leu Tyr Asn Tyr Trp
 35 40 45
 Phe Leu Tyr Ile Pro Tyr Leu Val Trp Phe Tyr Tyr Asp Trp Arg Thr
 50 55 60
 Pro Glu Gln Gly Gly Arg Arg Trp Asn Trp Val Gln Ser Trp Pro Val
 65 70 75 80
 Trp Lys Tyr Phe Lys Glu Tyr Phe Pro Ile Cys Leu Val Lys Thr Gln
 85 90 95
 Asp Leu Asp Pro Gly His Asn Tyr Ile Phe Gly Phe His Pro His Gly
 100 105 110
 Ile Phe Val Pro Gly Ala Phe Gly Asn Phe Cys Thr Lys Tyr Ser Asp
 115 120 125
 Phe Lys Lys Leu Phe Pro Gly Phe Thr Ser Tyr Leu His Val Ala Lys
 130 135 140
 Ile Trp Phe Cys Phe Pro Leu Phe Arg Glu Tyr Leu Met Ser Asn Gly
 145 150 155 160
 Pro Val Ser Val Ser Lys Glu Ser Leu Ser His Val Leu Ser Lys Asp
 165 170 175
 Gly Gly Gly Asn Val Ser Ile Ile Val Leu Gly Gly Ala Lys Glu Ala
 180 185 190
 Leu Glu Ala His Pro Gly Thr Phe Thr Leu Cys Ile Arg Gln Arg Lys
 195 200 205
 Gly Phe Val Lys Met Ala Leu Thr His Gly Ala Ser Leu Val Pro Val
 210 215 220
 Phe Ser Phe Gly Glu Asn Asp Leu Tyr Lys Gln Ile Asn Asn Pro Lys
 225 230 235 240

Gly Ser Trp Leu Arg Thr Ile Gln Asp Ala Met Tyr Asp Ser Met Gly
 245 250 255
 Val Ala Leu Pro Leu Ile Tyr Ala Arg Gly Ile Phe Gln His Tyr Phe
 260 265 270
 Gly Ile Met Pro Tyr Arg Lys Leu Ile Tyr Thr Val Val Gly Arg Pro
 275 280 285
 Ile Pro Val Gln Gln Ile Leu Asn Pro Thr Ser Glu Gln Ile Glu Glu
 290 295 300
 Leu His Gln Thr Tyr Leu Glu Glu Leu Lys Lys Leu Phe Asn Glu His
 305 310 315 320
 Lys Gly Lys Tyr Gly Ile Pro Glu His Glu Thr Leu Val Phe Lys
 325 330 335

<210> 7

<211> 1129

<212> DNA

<213> Homo sapiens

<400> 7

cgtgggtgca ggctgcagtg gctggcgccg tccctgcccc gccaggccat gaaggtagag 60
 ttgtcacccg tcaacatcca gctggcgccg cggctgcaga cgggtggccgt gctgcagtg 120
 gtccttttctt ttcttacagg gccgatgtcc attggaatca ctgtgatgct gatcatacac 180
 aactattttgt tcctttacat cccttatttg atgtggcttt actttgactg gcatacccca 240
 gagcgaggag gcaggagatc cagctggatc aaaaattgga ctctttggaa acactttaag 300
 gactattttc caattcatct tatcaaaact caagatttgg atccaagtca caactatata 360
 tttgggtttc acccccatgg aataatggca gttggagcct ttgggaattt ttctgtaaat 420
 tattctgact tcaaggacct gtttcctggc tttacttcat atcttcacgt gctgccactt 480
 tggttctggg gtcctgtctt tgcgagaatat gtgatgagtg ttgggtgggt ttccagtttcc 540
 aagaaaagtg tgtcctacat ggtaagcaag gaggggagtg gaaccatctc tgtcattgtc 600
 cttgggggtg caaaagaatc actggatgct catcctggaa agttcactct gttcatccgc 660
 cagcggaag gatttggtta aattgctttg acccatggcg cctctctgggt cccagtggtt 720
 tcttttggtg aaaatgaact gtttaacaa actgacaacc ctgaaggatc atggattaga 780
 actgttcaga ataaactgca gaagatcatg gggtttgctt tgcccctggt tcatgccagg 840
 ggagtttttc agtacaattt tggcctaatt acctatagga aagccatcca cactgttggt 900
 ggccgcccga tccctgttcg tcagactctg aaccgcaccc aggagcagat tgaggagtta 960
 catcagacct atatggagga acttaggaaa ttgtttgagg aacacaaaagg aaagtatggc 1020
 attccagagc acgagactct tgttttaaaa tgacttgact ataaaaaaaa attaaaaaat 1080
 aaaaataaat gacttggctg taataaggca taaagaagga taagagacc 1129

<210> 8

<211> 334

<212> PRT

<213> Homo sapiens

<400> 8

Met Lys Val Glu Phe Ala Pro Leu Asn Ile Gln Leu Ala Arg Arg Leu
 1 5 10 15
 Gln Thr Val Ala Val Leu Gln Trp Val Leu Ser Phe Leu Thr Gly Pro
 20 25 30
 Met Ser Ile Gly Ile Thr Val Met Leu Ile Ile His Asn Tyr Leu Phe
 35 40 45
 Leu Tyr Ile Pro Tyr Leu Met Trp Leu Tyr Phe Asp Trp His Thr Pro
 50 55 60
 Glu Arg Gly Gly Arg Arg Ser Ser Trp Ile Lys Asn Trp Thr Leu Trp
 65 70 75 80
 Lys His Phe Lys Asp Tyr Phe Pro Ile His Leu Ile Lys Thr Gln Asp
 85 90 95
 Leu Asp Pro Ser His Asn Tyr Ile Phe Gly Phe His Pro His Gly Ile
 100 105 110
 Met Ala Val Gly Ala Phe Gly Asn Phe Ser Val Asn Tyr Ser Asp Phe
 115 120 125
 Lys Asp Leu Phe Pro Gly Phe Thr Ser Tyr Leu His Val Leu Pro Leu
 130 135 140
 Trp Phe Trp Cys Pro Val Phe Arg Glu Tyr Val Met Ser Val Gly Leu
 145 150 155 160
 Val Ser Val Ser Lys Lys Ser Val Ser Tyr Met Val Ser Lys Glu Gly

<400>	10															
Met	Lys	Thr	Glu	His	Leu	Gln	Ser	Leu	Ser	Leu	Leu	Gln	Trp	Pro	Leu	
1				5					10					15		
Ser	Tyr	Val	Ala	Met	Phe	Trp	Ile	Val	Gln	Pro	Leu	Leu	Ile	Cys	Leu	
			20					25					30			
Leu	Phe	Thr	Pro	Leu	Trp	Pro	Leu	Pro	Thr	Val	Tyr	Phe	Val	Trp	Leu	
		35					40					45				
Leu	Leu	Asp	Trp	Lys	Thr	Pro	Asp	Lys	Gly	Gly	Arg	Arg	Ser	Asp	Trp	
	50					55					60					
Val	Arg	Asn	Trp	Asn	Val	Trp	Asn	His	Ile	Arg	Asp	Tyr	Phe	Pro	Ile	
65				70					75						80	
Thr	Ile	Leu	Lys	Thr	Lys	Asp	Leu	Ser	Pro	Ser	Glu	Asn	Tyr	Ile	Met	
			85						90					95		
Gly	Val	His	Pro	His	Gly	Leu	Leu	Thr	Phe	Gly	Ala	Phe	Cys	Asn	Phe	
			100					105					110			
Cys	Thr	Glu	Ala	Thr	Gly	Phe	Ser	Lys	Thr	Phe	Pro	Gly	Ile	Thr	Pro	
		115					120					125				
His	Leu	Ala	Thr	Leu	Ser	Trp	Phe	Phe	Lys	Ile	Pro	Ile	Ile	Arg	Asp	
	130					135						140				

Tyr Ile Met¹ Ala Lys Gly Leu Cys Ser Val Ser Gln Ala Ser Ile Asp
 145 150 155 160
 Tyr Leu Leu Ser His Gly Thr Gly Asn Leu Val Gly Ile Pro Ile Ile
 165 170 175
 Thr Val Val Gly Glu Ala Leu Pro Leu Pro Gln Val Lys Asn Pro Ser
 180 185 190
 Pro Glu Ile Val Asp Lys Tyr His Ala Leu Tyr Met Asp Ala Leu Tyr
 195 200 205
 Lys Leu Phe Glu Gln His Lys Val Gln Tyr Gly Cys Ser Asn Thr Gln
 210 215 220
 Lys Leu Ile Phe Leu
 225

<210> 11
 <211> 1240
 <212> DNA
 <213> Homo sapiens

<400> 11
 atcaactcag cttagaagt tttggccttc tggtaggct tcttgccaca acagaacagc 60
 accataacca tggctttctt ctcccgactg aatctccagg agggcctcca aaccttcttt 120
 gttttgcaat ggatcccagt ctatatattt ttagtttgga tcttgccagcc attgttcgtc 180
 tacctgctgt ttacatcctt gtggccgcta ccagtgcctt actttgcctg gttgttcctg 240
 gactggaaga cccagagcgg aggtggcagg cgcttcggcct gggtaaggaa ctgggtgtgtc 300
 tggacccaca tcaggggacta tttcccccatt acgacacctga agacaaagga cctatcacct 360
 gagcacaact acctcatggg gggttcacccc catggcctcc tgacqtttgg cgccttctgc 420
 aactttctgca ctgaggccac aggccttctcg aagaccttcc caggcatcac tcctcacttg 480
 gccacgctgt cctggttctt caagatcccc tttgttaggg agtacctcat ggccaaagg 540
 gtgtgctctg tgagccagcc agccatcaac tatctgctga gccatggcac tggcaacctc 600
 gtggggcattg tagtgggagg tgtgggtgag gccctgcaaa gtgtgcccaa caccaccacc 660
 ctcatcctcc agaagcgcaa ggggttcgtg cgcacagccc tccagcatgg ggcataacct 720
 gtcccttcat attccttttg tgagaacgaa gttttcaatc aggagacctt ccctgagggc 780
 acgtgggttaa gggtgttcca aaaaaccttc caggacacat tcaaaaaaat cctgggacta 840
 aatttctgta ccttccatgg ccggggcttc actcgcggat cctggggctt cctgcctttc 900
 aatcgcccca ttaccactgt tgttggggaa ccccttccaa ttcccaggat taaggagcca 960
 aaccagaaga cagtagacaa gtatcacgca ctctacatca gtgccctgcg caagctcttt 1020
 gaccaacaca aagttgaata tggcctccct gagaccgaag agctgacaat tacataacag 1080
 gagccacatt ccccatgtat caacccccaa agccatgagg gatccaagta gagccacaga 1140
 aaaagaagaa ttccaggaga gggaaagatc gtaaggatga gagaggagac catccaagcc 1200
 agaaattatt taataaatca gagttctagc aatagagtcc 1240

<210> 12
 <211> 335
 <212> PRT
 <213> Homo sapiens

<400> 12
 Met Ala Phe Phe Ser Arg Leu Asn Leu Gln Glu Gly Leu Gln Thr Phe
 1 5 10 15
 Phe Val Leu Gln Trp Ile Pro Val Tyr Ile Phe Leu Val Trp Ile Leu
 20 25 30
 Gln Pro Leu Phe Val Tyr Leu Leu Phe Thr Ser Leu Trp Pro Leu Pro
 35 40 45
 Val Leu Tyr Phe Ala Trp Leu Phe Leu Asp Trp Lys Thr Pro Glu Arg
 50 55 60
 Gly Gly Arg Arg Ser Ala Trp Val Arg Asn Trp Cys Val Trp Thr His
 65 70 75 80
 Ile Arg Asp Tyr Phe Pro Ile Thr Ile Leu Lys Thr Lys Asp Leu Ser
 85 90 95
 Pro Glu His Asn Tyr Leu Met Gly Val His Pro His Gly Leu Leu Thr
 100 105 110
 Phe Gly Ala Phe Cys Asn Phe Cys Thr Glu Ala Thr Gly Phe Ser Lys
 115 120 125
 Thr Phe Pro Gly Ile Thr Pro His Leu Ala Thr Leu Ser Trp Phe Phe
 130 135 140
 Lys Ile Pro Phe Val Arg Glu Tyr Leu Met Ala Lys Gly Val Cys Ser

145					150					155				160
Val	Ser	Gln	Pro	Ala	Ile	Asn	Tyr	Leu	Leu	Ser	His	Gly	Thr	Gly Asn
				165					170					175
Leu	Val	Gly	Ile	Val	Val	Gly	Gly	Val	Gly	Glu	Ala	Leu	Gln	Ser Val
			180					185					190	
Pro	Asn	Thr	Thr	Leu	Ile	Leu	Gln	Lys	Arg	Lys	Gly	Phe	Val	Arg
		195				200					205			
Thr	Ala	Leu	Gln	His	Gly	Ala	Tyr	Leu	Val	Pro	Ser	Tyr	Ser	Phe Gly
	210				215						220			
Glu	Asn	Glu	Val	Phe	Asn	Gln	Glu	Thr	Phe	Pro	Glu	Gly	Thr	Trp Leu
	225				230				235					240
Arg	Leu	Phe	Gln	Lys	Thr	Phe	Gln	Asp	Thr	Phe	Lys	Lys	Ile	Leu Gly
			245					250					255	
Leu	Asn	Phe	Cys	Thr	Phe	His	Gly	Arg	Gly	Phe	Thr	Arg	Gly	Ser Trp
		260					265					270		
Gly	Phe	Leu	Pro	Phe	Asn	Arg	Pro	Ile	Thr	Thr	Val	Val	Gly	Glu Pro
	275					280					285			
Leu	Pro	Ile	Pro	Arg	Ile	Lys	Arg	Pro	Asn	Gln	Lys	Thr	Val	Asp Lys
	290				295					300				
Tyr	His	Ala	Leu	Tyr	Ile	Ser	Ala	Leu	Arg	Lys	Leu	Phe	Asp	Gln His
	305				310				315					320
Lys	Val	Glu	Tyr	Gly	Leu	Pro	Glu	Thr	Gln	Glu	Leu	Thr	Ile	Thr
				325				330						335

<210> 13
 <211> 1872
 <212> DNA
 <213> Homo sapiens

<400> 13

aattcggctt	actcactata	gggctcgagc	ggcccccggg	caggtgccga	cttcatttcc	60
aagtctgcac	acaatgcagg	cagtagccat	gcctgacagc	cacatgacag	atactacacc	120
gctgaatgtg	ctctaaccct	ggacttggca	ttgccccctac	tgttgaggaa	gcagtgcgtt	180
tttctccagt	ctttcaggtc	ccttcaccag	ggaaccatta	acttggtgat	cagaacaagg	240
acatttccctt	acatttccctg	aaacacagtc	ctttcagttt	actctttttt	tgaggggggg	300
gcgcgggggaa	cggagtctcg	ctctgtcgcc	caggtctggag	tgcaatgggtg	caatctcagc	360
tcaactgcaac	ctctgcctcc	caggtcccaag	cgattctcct	gcctcagcct	cccgggtagc	420
cgggactaca	ggcgctgcc	accacgccc	gctaattttt	gtatttttag	tagagacgag	480
gtttcgcctg	gttggcaggc	tggtcttggg	actcctgacc	tcaggtgatt	tactcgctc	540
ggcctcccaa	agtgtctggg	ttacaggcat	gagccactgt	gcccagtcac	aagtttttat	600
tttagccatt	ttgataagtg	tgaagttccc	tgatggctaa	tgatgttcc	tttccatgt	660
gtcattttgt	catctatgcc	agagaagatt	tggagaggag	gacgtgaatt	ggaggaaaac	720
tgttccagga	ttccccacct	ctgggtggcc	accgctggct	cactgccatt	gaccacactg	780
caggcagagc	ctagtgcagt	gctggagcag	ggcccagaga	ggagagggt	tacagtgtga	840
attcagctca	gctggggaag	aagacacctt	cccttctaga	cctgaatcgg	gttcccaagc	900
aaccactgtg	attgctgtca	acctctacct	ggtggtgttc	acaccatact	ggcctgtcac	960
tgtgcttatt	cttacctggc	tggtttttga	ctggaagacc	cctcagcgag	gcggccgccg	1020
gtttacctgt	gtgaggcact	ggcgctgtg	gaaacactac	agcgattatt	tcctctctca	1080
gcttctgaag	actcatgaca	tctgccccag	ccgcaactac	atcctcgtct	gccaccctca	1140
tggtctcttt	gcccattgat	ggtttggcca	ctttgccaca	gaggcctcag	gcttctccaa	1200
gatatttccct	ggcatcaccc	cttacatact	cacactggga	gcctttttct	ggatgccttt	1260
cctcagagaa	tatgtaattg	ctacaggggc	ctgctctgtg	agtcgatact	ccattgactt	1320
tctgctgact	cataaaggca	caggcaacat	ggtcattgtg	gtgattgggtg	gactggctga	1380
gtgcagatac	agcctgccag	gttcttctac	cctgggtgtg	aagaaccggt	ctggctttgt	1440
gcgcattggc	cttcagcatg	gggtgctct	aatacctgcc	tatgctttg	gggagacgga	1500
cctctatgat	cagcacattt	tcactcctgg	tggctttgtc	aaccgcttcc	agaagtgtgt	1560
ccagagcatg	gtacacatct	accttgtgc	tttctatgga	cgtggcttca	ccaagaactc	1620
ctggggcctt	ctgccctata	gtcgccctgt	aaccaccatc	gtcggggagc	ctctaccaat	1680
gcccagatt	gagaatccaa	gccaggagat	cgtggctaaa	tatcacacac	tctatattga	1740
tgccctacgt	aaactgtttg	accagcataa	gaccaagtgt	ggtatctcag	agaccagga	1800
gctggagata	atttgacaga	catccccagt	agccttcacc	ctggctggaa	ggtatggatg	1860
gacccagtga	ga					1872

<210> 14
 <211> 333
 <212> PRT

<213> Homo sapiens

<400> 14

```

Met Leu Leu Pro Ser Lys Lys Asp Leu Lys Thr Ala Leu Asp Val Phe
 1      5      10      15
Ala Val Phe Gln Trp Ser Phe Ser Ala Leu Leu Ile Thr Thr Val
 20      25      30
Ile Ala Val Asn Leu Tyr Leu Val Val Phe Thr Pro Tyr Trp Pro Val
 35      40      45
Thr Val Leu Ile Leu Thr Trp Leu Ala Phe Asp Trp Lys Thr Pro Gln
 50      55      60
Arg Gly Gly Arg Arg Phe Thr Cys Val Arg His Trp Arg Leu Trp Lys
 65      70      75      80
His Tyr Ser Asp Tyr Phe Pro Leu Lys Leu Leu Lys Thr His Asp Ile
 85      90      95
Cys Pro Ser Arg Asn Tyr Ile Leu Val Cys His Pro His Gly Leu Phe
100      105      110
Ala His Gly Trp Phe Gly His Phe Ala Thr Glu Ala Ser Gly Phe Ser
115      120      125
Lys Ile Phe Pro Gly Ile Thr Pro Tyr Ile Leu Thr Leu Gly Ala Phe
130      135      140
Phe Trp Met Pro Phe Leu Arg Glu Tyr Val Met Ser Thr Gly Ala Cys
145      150      155      160
Ser Val Ser Arg Ser Ser Ile Asp Phe Leu Leu Thr His Lys Gly Thr
165      170      175
Gly Asn Met Val Ile Val Val Ile Gly Gly Leu Ala Glu Cys Arg Tyr
180      185      190
Ser Leu Pro Gly Ser Ser Thr Leu Val Leu Lys Asn Arg Ser Gly Phe
195      200      205
Val Arg Met Ala Leu Gln His Gly Val Pro Leu Ile Pro Ala Tyr Ala
210      215      220
Phe Gly Glu Thr Asp Leu Tyr Asp Gln His Ile Phe Thr Pro Gly Gly
225      230      235      240
Phe Val Asn Arg Phe Gln Lys Trp Phe Gln Ser Met Val His Ile Tyr
245      250      255
Pro Cys Ala Phe Tyr Gly Arg Gly Phe Thr Lys Asn Ser Trp Gly Leu
260      265      270
Leu Pro Tyr Ser Arg Pro Val Thr Thr Ile Val Gly Glu Pro Leu Pro
275      280      285
Met Pro Lys Ile Glu Asn Pro Ser Gln Glu Ile Val Ala Lys Tyr His
290      295      300
Thr Leu Tyr Ile Asp Ala Leu Arg Lys Leu Phe Asp Gln His Lys Thr
305      310      315      320
Lys Phe Gly Ile Ser Glu Thr Gln Glu Leu Glu Ile Ile
325      330

```

<210> 15

<211> 1050

<212> DNA

<213> Homo sapiens

<400> 15

```

ccacagcaga gctcacagaa cctgcgggag ccaggctgac ccgccagcat ggtagagttc      60
gccccttgt ttgtgccgtg ggagcgcagg ctgcagacac ttgctgtcct acagtttgtc      120
ttctccttct tggcactggg taagatctgc actgtgggct tcatagccct cctgtttaca      180
agattctggc tcctcactgt cctgtatgcg gcctgggtgt atctggaccg agacaagcca      240
cggcaggggg gcccgcacat ccaggccatc aggtgctgga ctatatggaa gtacatgaag      300
gactatttcc ccatccagct ggtcaagact gctgagctgg acccctctcg gaactacatt      360
gcgggcttcc acccccatgg agtcctggca gtcggagcct ttgccaacct gtgcactgag      420
agcacaggct tctcttcgat cttccccggg atccgcccc atctgatgat gctgaccttg      480
tggttccggg ccccttctct cagagattac atcatgtctg cagggttggg cacatcagaa      540
aaggagagtg ctgctcacat tctgaacagt aagggtggcg gaaacttgct gggcatcatt      600
gtagggggtg cccaggaggc cctggatgcc aggcctggat ccttcacgct gttactgcgg      660
aaccgaaagg gcttcgtcag gctcgccctg acacacgggg cacccttggg gccaatcttc      720
tccttcgggg agaatgacct atttgaccag attcccaact cttctggctc ctgggttaacg      780
tatatccaga atcggttgca gaagatcatg ggcactctcc tcccactctt tcatggccgt      840

```



```

gggtgtcttcc agtacagctt tggtttaata ccctaccgcc ggcccatcac cactgtgggg 900
aagcccatcg aggtacagaa gacgctgcat ccctcggagg aggaggtgaa ccagctgcac 960
cagcattata tcaaagagct gtgcaacctc ttcgaggccc acaaacttaa gttcaacatc 1020
cctgctgacc agcacttgga gttctgctga 1050

```

<210> 16
 <211> 333
 <212> PRT
 <213> Homo sapiens

```

<400> 16
Met Val Glu Phe Ala Pro Leu Phe Val Pro Trp Glu Arg Arg Leu Gln
1 5 10 15
Thr Leu Ala Val Leu Gln Phe Val Phe Ser Phe Leu Ala Leu Gly Lys
20 25 30
Ile Cys Thr Val Gly Phe Ile Ala Leu Leu Phe Thr Arg Phe Trp Leu
35 40 45
Leu Thr Val Leu Tyr Ala Ala Trp Trp Tyr Leu Asp Arg Asp Lys Pro
50 55 60
Arg Gln Gly Gly Arg His Ile Gln Ala Ile Arg Cys Trp Thr Ile Trp
65 70 75 80
Lys Tyr Met Lys Asp Tyr Phe Pro Ile Gln Leu Val Lys Thr Ala Glu
85 90 95
Leu Asp Pro Ser Arg Asn Tyr Ile Ala Gly Phe His Pro His Gly Val
100 105 110
Leu Ala Val Gly Ala Phe Ala Asn Leu Cys Thr Glu Ser Thr Gly Phe
115 120 125
Ser Ser Ile Phe Pro Gly Ile Arg Pro His Leu Met Met Leu Thr Leu
130 135 140
Trp Phe Arg Ala Pro Phe Phe Arg Asp Tyr Ile Met Ser Ala Gly Leu
145 150 155 160
Val Thr Ser Glu Lys Glu Ser Ala Ala His Ile Leu Asn Arg Lys Gly
165 170 175
Gly Gly Asn Leu Leu Gly Ile Ile Val Gly Gly Ala Gln Glu Ala Leu
180 185 190
Asp Ala Arg Pro Gly Ser Phe Thr Leu Leu Leu Arg Asn Arg Lys Gly
195 200 205
Phe Val Arg Leu Ala Leu Thr His Gly Ala Pro Leu Val Pro Ile Phe
210 215 220
Ser Phe Gly Glu Asn Asp Leu Phe Asp Gln Ile Pro Asn Ser Ser Gly
225 230 235 240
Ser Trp Leu Arg Tyr Ile Gln Asn Arg Leu Gln Lys Ile Met Gly Ile
245 250 255
Ser Leu Pro Leu Phe His Gly Arg Gly Val Phe Gln Tyr Ser Phe Gly
260 265 270
Leu Ile Pro Tyr Arg Arg Pro Ile Thr Thr Val Gly Lys Pro Ile Glu
275 280 285
Val Gln Lys Thr Leu His Pro Ser Glu Glu Glu Val Asn Gln Leu His
290 295 300
Gln His Tyr Ile Lys Glu Leu Cys Asn Leu Phe Glu Ala His Lys Leu
305 310 315 320
Lys Phe Asn Ile Pro Ala Asp Gln His Leu Glu Phe Cys
325 330

```

<210> 17
 <211> 10
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> synthetic peptide

```

<400> 17
Met Gly Asp Tyr Lys Asp Asp Asp Asp Gly
1 5 10

```

<210> 18
 <211> 1233
 <212> DNA
 <213> Homo sapiens

<400> 18
 ttcagccatg aagaccctca tagccgccta ctccgggggtc ctgcccggcg agcgtcaggc 60
 cgaggctgac cggagccagc gctctcacgg aggaacctgcg ctgtcgcgcg aggggtctgg 120
 gagatggggc actggatcca gcatcctctc cgcctccag gacctcttct ctgtcacctg 180
 gctcaatagg tccaaggtgg aaaagcagct acagggtcatc tcagtgtctc agtgggtcct 240
 gtcttctctt gtactgggag tggcctgcag tgccatcctc atgtacatat tctgcactga 300
 ttgtgtggctc atcgctgtgc tctacttcac ttggctgggtg tttgactgga acacacccaa 360
 gaaaggtggc aggaggtcac agtgggtccg aaactgggct gtgtggcgct actttcgaga 420
 ctactttccc atccagctgg tgaagacaca caacctgctg accaccagga actatatctt 480
 tggataccac ccccatggta tcatgggcct ggggtgccttc tgcaacttca gcacagaggc 540
 cacagaagtg agcaagaagt tcccaggcat acggccttac ctggctacac tggcaggcaa 600
 cttccgaatg cctgtgttga gggagtacct gatgtctgga ggtatctgcc ctgtcagccg 660
 ggacaccata gactatttgc tttcaaagaa tgggagtggc aatgctatca tcatcgtggg 720
 cgggggtgcg gctgagctct tgagctccat gcctggcaag aatgcagtca ccctgcccga 780
 ccgcaagggc tttgtgaaac tggccctgcg tcatggagct gacctgggtc ccatctactc 840
 ctttggagag aatgaagtgt acaagcaggt gatcttcgag gagggctcct ggggcccagtg 900
 ggtccagaag aagttccaga aatacattgg tttcgcacca tgcattctcc atggtcgagg 960
 cctcttctcc tccgacacct gggggctggg gccctactcc aagcccatca ccaactgttg 1020
 gggagagccc atcaccatcc ccaagctgga gcacccaacc cagcaagaca tcgacctgta 1080
 ccacaccatg tacatggagg ccctggtgaa gctcttcgac aagcacaaga ccaagttcgg 1140
 cctcccggag actgaggtcc tggaggtgaa ctgagccagc cttcggggcc aattccctgg 1200
 aggaaccagc tgcaaatac ttttttctc tgt 1233

<210> 19
 <211> 284
 <212> PRT
 <213> Homo sapiens

<400> 19
 Met Val Glu Phe Ala Pro Leu Phe Met Pro Trp Glu Arg Arg Leu Gln
 1 5 10 15
 Thr Leu Ala Val Leu Gln Phe Val Phe Ser Phe Leu Ala Leu Ala Glu
 20 25 30
 Ile Cys Thr Val Gly Phe Ile Ala Leu Leu Phe Thr Arg Phe Trp Leu
 35 40 45
 Leu Thr Val Leu Tyr Ala Ala Trp Trp Tyr Leu Asp Arg Asp Lys Pro
 50 55 60
 Arg Gln Gly Gly Arg His Ile Gln Ala Ile Arg Cys Trp Thr Ile Trp
 65 70 75 80
 Lys Tyr Met Lys Asp Tyr Phe Pro Ile Ser Leu Val Lys Thr Ala Glu
 85 90 95
 Leu Asp Pro Ser Arg Asn Tyr Ile Ala Gly Phe His Pro His Gly Val
 100 105 110
 Leu Ala Val Gly Ala Phe Ala Asn Leu Cys Thr Glu Ser Thr Gly Phe
 115 120 125
 Ser Ser Ile Phe Pro Gly Ile Arg Pro His Leu Met Met Leu Thr Leu
 130 135 140
 Trp Phe Arg Ala Pro Phe Phe Arg Asp Tyr Ile Met Ser Ala Gly Leu
 145 150 155 160
 Val Thr Ser Glu Lys Glu Ser Ala Ala His Ile Leu Asn Arg Lys Gly
 165 170 175
 Gly Gly Asn Leu Leu Gly Ile Ile Val Gly Gly Ala Gln Glu Ala Leu
 180 185 190
 Asp Ala Arg Pro Gly Ser Phe Thr Leu Leu Leu Arg Asn Arg Lys Gly
 195 200 205
 Phe Val Arg Leu Ala Leu Thr His Gly Tyr Gln Ala Ser Gly Lys Ser
 210 215 220
 Thr Leu Gly Ser Val Gly Asn Trp Gln Gly Phe Tyr Phe Gly Gly Lys
 225 230 235 240
 Met Ala Glu Thr Asn Ala Asp Ser Ile Leu Val Glu Ile Phe Ser Pro
 245 250 255

Phe Thr Ile Lys Ile Ile Phe Trp Cys Leu Met Pro Lys Tyr Leu Glu
 260 265 270
 Lys Phe Pro Gln Arg Arg Leu Ser Asp Leu Arg Asn
 275 280

<210> 20
 <211> 334
 <212> PRT
 <213> Homo sapiens

<400> 20
 Met Val Glu Phe Ala Pro Leu Phe Val Pro Trp Glu Arg Arg Leu Gln
 1 5 10 15
 Thr Leu Ala Val Leu Gln Phe Val Phe Ser Phe Leu Ala Leu Ala Glu
 20 25 30
 Ile Cys Thr Val Gly Phe Ile Ala Leu Leu Phe Thr Arg Phe Trp Leu
 35 40 45
 Leu Thr Val Leu Tyr Ala Ala Trp Trp Tyr Leu Asp Arg Asp Lys Pro
 50 55 60
 Arg Gln Gly Gly Arg His Ile Gln Ala Ile Arg Cys Trp Thr Ile Trp
 65 70 75 80
 Lys Tyr Met Lys Asp Tyr Phe Pro Ile Ser Leu Val Lys Thr Ala Glu
 85 90 95
 Leu Asp Pro Ser Arg Asn Tyr Ile Ala Gly Phe His Pro His Gly Val
 100 105 110
 Leu Ala Val Gly Ala Phe Ala Asn Leu Cys Thr Glu Ser Thr Gly Phe
 115 120 125
 Ser Ser Ile Phe Pro Gly Ile Arg Pro His Leu Met Met Leu Thr Leu
 130 135 140
 Trp Phe Arg Ala Pro Phe Phe Arg Asp Tyr Ile Met Ser Ala Gly Leu
 145 150 155 160
 Val Thr Ser Glu Lys Glu Ser Ala Ala His Ile Leu Asn Arg Lys Gly
 165 170 175
 Gly Gly Asn Leu Leu Gly Ile Ile Val Gly Gly Ala Gln Glu Ala Leu
 180 185 190
 Asp Ala Arg Pro Gly Ser Phe Thr Leu Leu Leu Arg Asn Arg Lys Gly
 195 200 205
 Phe Val Arg Leu Ala Leu Thr His Gly Ala Pro Leu Val Pro Ile Phe
 210 215 220
 Ser Phe Gly Glu Asn Asp Leu Phe Asp Gln Ile Pro Asn Ser Ser Gly
 225 230 235 240
 Ser Trp Leu Arg Tyr Ile Gln Asn Arg Leu Gln Lys Ile Met Gly Ile
 245 250 255
 Ser Leu Pro Leu Phe His Gly Arg Gly Val Phe Gln Tyr Ser Phe Gly
 260 265 270
 Leu Ile Pro Tyr Arg Arg Pro Ile Thr Thr Val Val Gly Lys Pro Ile
 275 280 285
 Glu Val Gln Lys Thr Leu His Pro Ser Glu Glu Glu Val Asn Gln Leu
 290 295 300
 His Gln Arg Tyr Ile Lys Glu Leu Cys Asn Leu Phe Glu Ala His Lys
 305 310 315 320
 Leu Lys Phe Asn Ile Pro Ala Asp Gln His Leu Glu Phe Cys
 325 330

<210> 21
 <211> 296
 <212> PRT
 <213> Mus musculus

<400> 21
 Met Val Glu Phe Ala Pro Leu Leu Val Pro Trp Glu Arg Arg Leu Gln
 1 5 10 15
 Thr Phe Ala Val Leu Gln Trp Val Phe Ser Phe Leu Ala Leu Ala Gln
 20 25 30
 Leu Cys Ile Val Ile Phe Val Gly Leu Leu Phe Thr Arg Phe Trp Leu
 35 40 45

Phe Ser Val Leu Tyr Ala Thr Trp Trp Tyr Leu Asp Trp Asp Lys Pro
 50 55 60
 Arg Gln Gly Gly Arg Pro Ile Gln Phe Phe Arg Arg Leu Ala Ile Trp
 65 70 75 80
 Lys Tyr Met Lys Asp Tyr Phe Pro Val Ser Leu Val Lys Thr Ala Glu
 85 90 95
 Leu Asp Pro Ser Arg Asn Tyr Ile Ala Gly Phe His Pro His Gly Val
 100 105 110
 Leu Ala Ala Gly Ala Phe Leu Asn Leu Cys Thr Glu Ser Thr Gly Phe
 115 120 125
 Thr Ser Leu Phe Pro Gly Ile Arg Ser Tyr Leu Met Met Leu Thr Val
 130 135 140
 Trp Phe Arg Ala Pro Phe Phe Arg Asp Tyr Ile Met Ser Gly Gly Leu
 145 150 155 160
 Val Ser Ser Glu Lys Val Ser Ala Asp His Ile Leu Ser Arg Lys Gly
 165 170 175
 Gly Gly Asn Leu Leu Ala Ile Ile Val Gly Gly Ala Gln Glu Ala Leu
 180 185 190
 Asp Ala Arg Pro Gly Ala Tyr Arg Leu Leu Leu Lys Asn Arg Lys Gly
 195 200 205
 Phe Ile Arg Leu Ala Leu Met His Gly Ala Ala Leu Val Pro Ile Phe
 210 215 220
 Ser Phe Gly Glu Asn Asn Leu Phe Asn Gln Val Glu Asn Thr Pro Gly
 225 230 235 240
 Thr Trp Leu Arg Trp Ile Gln Asn Arg Leu Gln Lys Ile Met Gly Ile
 245 250 255
 Ser Leu Pro Leu Phe His Gly Arg Gly Val Phe Gln Tyr Ser Phe Gly
 260 265 270
 Leu Met Pro Phe Arg Gln Pro Ile Thr Thr Ile Gly Glu Arg Arg Ser
 275 280 285
 Arg Arg Ala Glu Gln Asp Gln Glu
 290 295

<210> 22

<211> 323

<212> PRT

<213> Mus musculus

<400> 22

Met Val Glu Phe Ala Pro Leu Leu Val Pro Trp Glu Arg Arg Leu Gln
 1 5 10 15
 Thr Phe Ala Val Leu Gln Trp Val Phe Ser Phe Leu Ala Leu Ala Gln
 20 25 30
 Leu Cys Ile Val Ile Phe Val Gly Leu Leu Phe Thr Arg Phe Trp Leu
 35 40 45
 Phe Ser Val Leu Tyr Ala Thr Trp Trp Tyr Leu Asp Trp Asp Lys Pro
 50 55 60
 Arg Gln Gly Gly Arg Pro Ile Gln Phe Phe Arg Arg Leu Ala Ile Trp
 65 70 75 80
 Lys Tyr Met Lys Asp Tyr Phe Pro Val Ser Leu Val Lys Thr Ala Glu
 85 90 95
 Leu Asp Pro Ser Arg Asn Tyr Ile Ala Gly Phe His Pro His Gly Val
 100 105 110
 Leu Ala Ala Gly Ala Phe Leu Asn Leu Cys Thr Glu Ser Thr Gly Phe
 115 120 125
 Thr Ser Leu Phe Pro Gly Ile Arg Ser Tyr Leu Met Met Leu Thr Val
 130 135 140
 Trp Phe Arg Ala Pro Phe Phe Arg Asp Tyr Ile Met Ser Gly Gly Leu
 145 150 155 160
 Val Ser Ser Glu Lys Val Ser Ala Asp His Ile Leu Ser Arg Lys Gly
 165 170 175
 Gly Gly Asn Leu Leu Ala Ile Ile Val Gly Gly Ala Gln Glu Ala Leu
 180 185 190
 Asp Ala Arg Pro Gly Ala Tyr Arg Leu Leu Leu Lys Asn Arg Lys Gly
 195 200 205
 Phe Ile Met Leu Ala Leu Met His Gly Ala Ala Leu Cys Ala Ile Phe

210	215	220
Ser Phe Gly Glu Asn Asn	Leu Phe Asn Gln Val	Glu Asn Thr Pro Gly
225	230	235
Thr Trp Leu Arg Trp Ile	Gln Asn Arg Leu Gln	Lys Ile Met Gly Ile
245	250	255
Ser Leu Pro Leu Phe His	Gly Arg Gly Val Phe	Gln Tyr Ser Phe Gly
260	265	270
Leu Met Pro Phe Arg Gln	Pro Ile Thr Thr Ile	Gly Glu Arg Arg Ser
275	280	285
Arg Pro Gly Arg Ala Gly	Ser Gly Ile Gly Pro	Glu Leu Asp Ala Thr
290	295	300
Arg Glu Leu Asp Asp Gly	Gln Glu Met Phe Pro	Ser Ser Arg Asn Ser
305	310	315
Gln Ser Lys		320

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 September 2002 (06.09.2002)

PCT

(10) International Publication Number
WO 02/068595 A3

- (51) International Patent Classification⁷: C12N 9/10, 9/00, 15/00, 1/12, C07H 21/04
- (21) International Application Number: PCT/US02/05474
- (22) International Filing Date: 21 February 2002 (21.02.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/271,307 23 February 2001 (23.02.2001) US
09/794,715 26 February 2001 (26.02.2001) US
10/046,924 14 January 2002 (14.01.2002) US
- (71) Applicant (for all designated States except US): **THE REGENTS OF THE UNIVERSITY OF CALIFORNIA** [US/US]; 1111 Franklin Street, 5th Floor, Oakland, CA 94607 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **CASES, Sylvaine** [FR/US]; 9 Meadow Park, Belmont, CA 94002 (US). **STONE, Scot** [CA/US]; 2550 Hilborn Dr., #168, Fairfield, CA 94533 (US). **ZHOU, Ping** [CN/US]; 886 Stonehaven Drive, Walnut Creek, CA 94598 (US). **FARESE, Robert, V.** [US/US]; 25 6th Avenue, San Francisco, CA 94118 (US). **YEN, Chi-Liang, Eric** [CN/US]; 1427 22nd Avenue, San Francisco, CA 94122 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- (88) Date of publication of the international search report:
31 October 2002
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: MONO-AND DIACYGLYCEROL ACYLTRANSFERASES AND METHODS OF USE THEREOF

(57) Abstract: Nucleic acid compositions encoding polypeptide products with diglyceride acyltransferase and/or monoacylglycerol acyltransferase activity, as well as the polypeptide products encoded thereby, i.e., mammalian DGAT2 and MGAT1 polypeptide products, and methods for producing the same, are provided. Also provided are: methods and compositions for modulating DGAT2 and MGAT1 activity; DGAT2 and MGAT1 transgenic cells, animals and plants, as well as methods for their preparation; and methods for making diglyceride, diglyceride compositions, triglycerides and triglyceride compositions, as well as the compositions produced by these methods. The subject methods and compositions find use in a variety of different applications, including research, medicine, agriculture and industry applications.

WO 02/068595 A3

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/05474

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C12N 9/10, 9/00, 15/00, 1/12; C07H 21/04

US CL : 435/193, 183, 320.1, 252.1; 536/23.1, 23.2, 23.5

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/193, 183, 320.1, 252.1; 536/23.1, 23.2, 23.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00/12708 A2 (GENENTECH INC.) 09 March 2000 (09.03.00), page 24 and Figures 163 and 164.	1, 2, 7-10
A	OELKERS et al. Characterization of Two Human Genes Encoding Acyl Coenzyme A: Cholesterol Acyltransferase-related Enzymes, Journal of Biological Chemistry. October 1998, Vol. 273. No. 41, pages 26765-26771.	1-3, 7-10
A	CASES et al. Identification of a Gene Encoding an Acyl CoA:diacylglycerol acyltransferase, a Key Enzyme in Triacylglycerol synthesis, Proceedings of the National Academy of Sciences, October 1998, Vol. 95, pages 13018-13023.	1-3, 7-10

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

18 July 2002 (18.07.2002)

Date of mailing of the international search report

28 AUG 2002

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Ponnathapu Achutamurthy

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/05474

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claim Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claim Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:
Please See Continuation Sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.: 1-3 and 7-10

Remark on Protest

☐
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/05474

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING

This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Groups I-IX, claim(s) 1-3, 7-10, drawn to a polynucleotide encoding a mono or diacylglycerol transferase, vectors and host cells comprising said polynucleotide and methods of expressing said polynucleotide, corresponding to SEQ ID NOs: 01, 03, 05, 07, 09, 11, 13, 15, and 18.

Groups X-XVIII, claim(s) 4-6, drawn to a mono or diacylglycerol transferase polypeptide encoded by SEQ ID NOs: 01, 03, 05, 07, 09, 11, 13, 15, and 18..

Groups XIX-XXVII, claim(s) 11-14, drawn to an antibody which binds to a mono or diacylglycerol transferase polypeptide encoded by SEQ ID NOs: 01, 03, 05, 07, 09, 11, 13, 15, and 18

Groups XXVIII-XXXVI, claim(s) 15-18, drawn to a method for inhibiting the activity of a mono or diacylglycerol transferase polypeptide encoded by SEQ ID NOs: 01, 03, 05, 07, 09, 11, 13, 15, and 18.

Group XXXVII, claim(s) 19-21, drawn to method of modulating a symptom in a mammalian host of a disease condition associated with the acyltransferase activity of a mono or diacylglycerol transferase polypeptide.

Group XXXVIII, claim(s) 22, drawn to method of producing a triacylglycerol.

Group XXXIX, claim(s) 23, drawn to method of identifying an agent that inhibits an acyltransferase activity of a mono or diacylglycerol transferase polypeptide.

The inventions listed as Groups I through XXXIX do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The technical feature of Groups I through XXXIX is a polynucleotide encoding a mono or diacylglycerol transferase polypeptide and GENENTECH INC. (WO 00/12708, March 9, 2000) teaches a 1570 polynucleotide sequence which is 98.8% identical to instantly disclosed SEQ ID NO: 1 and encodes a diacyltransferase protein. Therefore the shared technical feature of each of the groups is not a special technical feature. Further the different SEQ ID NOs have no common core structure. Further, inventions XXVIII through XXXIX are drawn to different methods which are not the same such that the method of Groups XXVIII-XXXVI comprise contacting a protein with an inhibitor of acyltransferase activity, the method of Group XXXVII comprises administering a pharmaceutical composition comprising a modulating agent, the method of Group XXXVIII comprises contacting a diacylglycerol and fatty acyl CoA with a diacylglycerol transferase polypeptide and the method of Group XXXIX comprises contacting a mono or diacylglycerol transferase polypeptide with a test agent.

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 September 2002 (06.09.2002)

PCT

(10) International Publication Number
WO 02/068595 A3

(51) International Patent Classification⁷: C12N 9/10,
9/00, 15/00, 1/12, C07H 21/04

(21) International Application Number: PCT/US02/05474

(22) International Filing Date: 21 February 2002 (21.02.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/271,307 23 February 2001 (23.02.2001) US
09/794,715 26 February 2001 (26.02.2001) US
10/046,924 14 January 2002 (14.01.2002) US

(71) Applicant (for all designated States except US): **THE REGENTS OF THE UNIVERSITY OF CALIFORNIA** [US/US]; 1111 Franklin Street, 5th Floor, Oakland, CA 94607 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CASES, Sylvaine** [FR/US]; 9 Meadow Park, Belmont, CA 94002 (US). **STONE, Scot** [CA/US]; 2550 Hilborn Dr., #168, Fairfield, CA 94533 (US). **ZHOU, Ping** [CN/US]; 886 Stonehaven Drive, Walnut Creek, CA 94598 (US). **FARESE, Robert, V.** [US/US]; 25 6th Avenue, San Francisco, CA 94118 (US). **YEN, Chi-Liang, Eric** [CN/US]; 1427 22nd Avenue, San Francisco, CA 94122 (US).

(74) Agent: **BORDEN, Paula, A.**; **BOZICEVIC, FIELD & FRANCIS, L.L.P.**, 200 Middlefield Road, Suite 200, Menlo Park, CA 94025 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
31 October 2002

(48) Date of publication of this corrected version:
24 December 2003

(15) Information about Correction:
see PCT Gazette No. 52/2003 of 24 December 2003, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: MONO-AND DIACYLGLYCEROL ACYLTRANSFERASES AND METHODS OF USE THEREOF

(57) Abstract: Nucleic acid compositions encoding polypeptide products with diglyceride acyltransferase and/or monoacylglycerol acyltransferase activity, as well as the polypeptide products encoded thereby, i.e., mammalian DGAT2 and MGAT1 polypeptide products, and methods for producing the same, are provided. Also provided are: methods and compositions for modulating DGAT2 and MGAT1 activity; DGAT2 and MGAT1 transgenic cells, animals and plants, as well as methods for their preparation; and methods for making diglyceride, diglyceride compositions, triglycerides and triglyceride compositions, as well as the compositions produced by these methods. The subject methods and compositions find use in a variety of different applications, including research, medicine, agriculture and industry applications.

WO 02/068595 A3